

Datasheet for ABIN7601207
anti-HGS antibody (AA 3-777)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	HGS
Binding Specificity:	AA 3-777
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This HGS antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Flow Cytometry (FACS)

Product Details

Purpose:	Anti-HGS Antibody Picoband® (monoclonal, 6C2E1)
Immunogen:	E.coli-derived human HGS recombinant protein (Position: R3-D777).
Clone:	6C2E1
Isotype:	IgG1
Cross-Reactivity (Details):	No cross-reactivity with other proteins.
Characteristics:	Anti-HGS Antibody Picoband® (monoclonal, 6C2E1) (ABIN7601207). Tested in Flow Cytometry, IHC, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.

Product Details

Purification: Immunogen affinity purified.

Target Details

Target: HGS

Alternative Name: HGS ([HGS Products](#))

Background: Synonyms: Bcl-2 homologous antagonist/killer, Apoptosis regulator BAK, Bcl-2-like protein 7, Bcl2-L-7, BAK1, BAK, BCL2L7, CDN1

Tissue Specificity: Expressed in a wide variety of tissues, with highest levels in the heart and skeletal muscle.

Background: Hepatocyte growth factor-regulated tyrosine kinase substrate is an enzyme that in humans is encoded by the HGS gene. It is mapped to 17q25.3. The protein encoded by this gene regulates endosomal sorting and plays a critical role in the recycling and degradation of membrane receptors. The encoded protein sorts monoubiquitinated membrane proteins into the multivesicular body, targeting these proteins for lysosome-dependent degradation.

Molecular Weight: 110 kDa

Gene ID: 9146

UniProt: [O14964](#)

Pathways: [EGFR Signaling Pathway](#), [CXCR4-mediated Signaling Events](#), [Synaptic Vesicle Exocytosis](#), [EGFR Downregulation](#)

Application Details

Application Notes: Western blot, 0.25-0.5 µg/mL, Human, Mouse, Rat

Immunohistochemistry(Paraffin-embedded Section), 2-5 µg/mL, Human

Flow Cytometry (Fixed), 1-3 µg/1x10⁶ cells, Human

1. Asao, H., Sasaki, Y., Arita, T., Tanaka, N., Endo, K., Kasai, H., Takeshita, T., Endo, Y., Fujita, T., Sugamura, K. Hrs is associated with STAM, a signal-transducing adaptor molecule: its suppressive effect on cytokine-induced cell growth. J. Biol. Chem. 272: 32785-32791, 1997. 2. Gutmann, D. H., Haipek, C. A., Burke, S. P., Sun, C.-X., Scoles, D. R., Pulst, S. M. The NF2 interactor, hepatocyte growth factor-regulated tyrosine kinase substrate (HRS), associates with merlin in the 'open' conformation and suppresses cell growth and motility. Hum. Molec. Genet. 10: 825-834, 2001. 3. Komada, M., Masaki, R., Yamamoto, A., Kitamura, N. Hrs, a tyrosine kinase substrate with a conserved double zinc finger domain, is localized to the cytoplasmic surface of early endosomes. J. Biol. Chem. 272: 20538-20544, 1997.

Application Details

Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Adding 0.2 mL of distilled water will yield a concentration of 500 µg/mL.
Concentration:	500 µg/mL
Buffer:	Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na2HPO4.
Storage:	4 °C,-20 °C
Storage Comment:	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.