

Datasheet for ABIN7601245  
**anti-LRRC47 antibody (AA 309-583)**



[Go to Product page](#)

## Overview

Quantity:	100 µg
Target:	LRRC47
Binding Specificity:	AA 309-583
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LRRC47 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Flow Cytometry (FACS)

## Product Details

Purpose:	Anti-LRRC47 Antibody Picoband®
Immunogen:	E.coli-derived human LRRC47 recombinant protein (Position: L309-R583).
Isotype:	IgG
Cross-Reactivity (Details):	No cross reactivity with other proteins.
Characteristics:	Anti-LRRC47 Antibody Picoband® (ABIN7601245). Tested in ELISA, WB, Flow Cytometry applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Purification:	Immunogen affinity purified.

## Target Details

Target:	LRRC47
Alternative Name:	LRRC47 ( <a href="#">LRRC47 Products</a> )
Background:	<p>Synonyms: 70 kDa ribosomal protein S6 kinase 1 antibody, KS6B1_HUMAN antibody, p70 alpha antibody, P70 beta 1 antibody, p70 ribosomal S6 kinase alpha antibody, p70 ribosomal S6 kinase beta 1 antibody, p70 S6 kinase alpha antibody, P70 S6 Kinase antibody, p70 S6 kinase alpha 1 antibody, p70 S6 kinase alpha 2 antibody, p70 S6K antibody, p70 S6K-alpha antibody, p70 S6KA antibody, p70(S6K) alpha antibody, p70(S6K)-alpha antibody, p70-alpha antibody, p70-S6K 1 antibody, p70-S6K antibody, P70S6K antibody, P70S6K1 antibody, p70S6Kb antibody, PS6K antibody, Ribosomal protein S6 kinase 70 kDa polypeptide 1 antibody, Ribosomal protein S6 kinase beta 1 antibody, Ribosomal protein S6 kinase beta-1 antibody, Ribosomal protein S6 kinase I antibody, RPS6KB1 antibody, S6K antibody, S6K-beta-1 antibody, S6K1 antibody, Serine/threonine kinase 14 alpha antibody, Serine/threonine-protein kinase 14A antibody, STK14A antibody</p> <p>Tissue Specificity: Expressed in all tissues.</p> <p>Background: Chromosome 1 is the largest human chromosome spanning about 260 million base pairs and making up 8 % of the human genome. There are about 3,000 genes on chromosome 1, and considering the great number of genes there are also a large number of diseases associated with chromosome 1. Notably, the rare aging disease Hutchinson-Gilford progeria is associated with the LMNA gene which encodes Lamin A. When defective, the LMNA gene product can build up in the nucleus and cause characteristic nuclear blebs. The mechanism of rapidly enhanced aging is unclear and is a topic of continuing exploration. The MUTYH gene is located on chromosome 1 and is partially responsible for familial adenomatous polyposis. Stickler syndrome, Parkinsons, Gaucher disease and Usher syndrome are also associated with chromosome 1. A breakpoint has been identified in 1q which disrupts the DISC1 gene and is linked to schizophrenia. Aberrations in chromosome 1 are found in a variety of cancers including head and neck cancer, malignant melanoma and multiple myeloma.</p>
Molecular Weight:	68 kDa
Gene ID:	57470
UniProt:	<a href="#">Q8N1G4</a>

## Application Details

Application Notes:	Western blot, 0.1-0.25 µg/mL, Human
	Flow Cytometry (Fixed), 1-3 µg/1x10 <sup>6</sup> cells, Human

## Application Details

---

ELISA, 0.1-0.5 µg/mL, -

1. Gross, M. B. Personal Communication. Baltimore, Md. 1/14/2021. 2. Hirosawa, M., Nagase, T., Ishikawa, K., Kikuno, R., Nomura, N., Ohara, O. Characterization of cDNA clones selected by the GeneMark analysis from size-fractionated cDNA libraries from human brain. DNA Res. 6: 329-336, 1999.

---

Restrictions: For Research Use only

## Handling

---

Format: Lyophilized

Reconstitution: Adding 0.2 mL of distilled water will yield a concentration of 500 µg/mL.

Concentration: 500 µg/mL

Buffer: Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na<sub>2</sub>HPO<sub>4</sub>.

Storage: 4 °C, -20 °C

Storage Comment: At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month.  
It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.