

Datasheet for ABIN7601365
anti-TNFRSF1A antibody (AA 336-440)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	TNFRSF1A
Binding Specificity:	AA 336-440
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TNFRSF1A antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Flow Cytometry (FACS)

Product Details

Purpose:	Anti-TNF Receptor I/TNFRSF1A Antibody Picoband®
Immunogen:	E.coli-derived human TNF Receptor I/TNFRSF1A recombinant protein (Position: N336-A440).
Isotype:	IgG
Cross-Reactivity (Details):	No cross-reactivity with other proteins.
Characteristics:	Anti-TNF Receptor I/TNFRSF1A Antibody Picoband® (ABIN7601365). Tested in ELISA, Flow Cytometry, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Purification:	Immunogen affinity purified.

Target Details

Target:	TNFRSF1A
Alternative Name:	TNFRSF1A (TNFRSF1A Products)
Background:	Synonyms: Tumor necrosis factor receptor superfamily member 1A, Tumor necrosis factor receptor 1, TNF-R1, Tumor necrosis factor receptor type I, TNF-RI, TNFR-I, p55, p60, CD120a, Tumor necrosis factor receptor superfamily member 1A, membrane form, Tumor necrosis factor-binding protein 1, TBPI, TNFRSF1A, TNFAR, TNFR1 Tissue Specificity: Activated T-cells. Highly expressed on tonsillar T-cells, which are closely associated with B-cells in the apical light zone of germinal centers, the site of terminal B- cell maturation. Expressed at lower levels in thymus, lung, lymph node and peripheral blood leukocytes. Expressed in the medulla of fetal and newborn thymus. Background: Tumor necrosis factor receptor superfamily member 1A (TNFRSF1A), also known as TNFR1, is a protein that in humans is encoded by the TNFRSF1A gene. The protein encoded by this gene is a member of the Tumor necrosis factor receptor superfamily, which also contains TNFRSF1B. The TNFR1 gene is mapped to 12pter-cen. It encodes a protein of 455 amino acids. And this receptor can activate the transcription factor NF-κB, mediate apoptosis, and function as a regulator of inflammation.
Molecular Weight:	60-70 kDa
Gene ID:	7132
UniProt:	P19438
Pathways:	NF-kappaB Signaling , Apoptosis , Caspase Cascade in Apoptosis , Hepatitis C , Ubiquitin Proteasome Pathway

Application Details

Application Notes:	Western blot, 0.25-0.5 µg/mL, Human, Mouse, Rat Flow Cytometry (Fixed), 1-3 µg/1×10⁶ cells, Human ELISA, 0.1-0.5 µg/mL, - 1. Baker E, Chen LZ, Smith CA, Callen DF, Goodwin R, Sutherland GR (Nov 1991). "Chromosomal location of the human tumor necrosis factor receptor genes". Cytogenet Cell Genet 57 (2-3): 117-8. 2. Schall TJ, Lewis M, Koller KJ, Lee A, Rice GC, Wong GH, Gatanaga T, Granger GA, Lentz R, Raab H, et al. (Jun 1990). "Molecular cloning and expression of a receptor for human tumor necrosis factor". Cell 61 (2): 361-70.
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Adding 0.2 mL of distilled water will yield a concentration of 500 µg/mL.
Concentration:	500 µg/mL
Buffer:	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage:	4 °C, -20 °C
Storage Comment:	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.