# antibodies -online.com





# Datasheet for ABIN760156

# anti-SERINC3 antibody (AA 51-150) (Cy5.5)



#### Overview

| Quantity:            | 100 μL   |
|----------------------|--|
| Target:              | SERINC3  |
| Binding Specificity: | AA 51-150  |
| Reactivity:          | Mouse  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This SERINC3 antibody is conjugated to Cy5.5   |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human SERINC3 |
|-----------------------|---|
| Isotype:              | IgG   |
| Cross-Reactivity:     | Mouse   |
| Predicted Reactivity: | Human,Rat,Dog,Cow,Horse,Chicken,Rabbit                      |
| Purification:         | Purified by Protein A.                                      |

## **Target Details**

| Target:           | SERINC3                    |
|-------------------|----------------------------|
| Alternative Name: | SERINC3 (SERINC3 Products) |

## **Target Details**

| Background:         | Synonyms: TMS 1, TMS-1, AIGP1, DF33, SBBI99, serine incorporator 3, TDE, TDE1, Tumor          |
|---------------------|---|
|                     | dferentially expressed protein 1, SERC3_HUMAN.  |
|                     | Background: TDE1, tumor differentially expressed protein 1, is a multi-pass membrane protein  |
|                     | also known as serine incorporator 3. Also found on the membranes of the Golgi apparatus       |
|                     | within cells, TDE1 is highly expressed in neuronal populations but is also found in thymus,   |
|                     | kidney, liver and testis. Expression levels of TDE1 in tumors can be as much as tenfold the   |
|                     | amount found in normal tissue of the same type. This increased expression implicates TDE1 as  |
|                     | being involved in the cellular transformation from normal to malignant tissue. It is believed |
|                     | TDE1 contributes to oncogenesis by partially protecting cells from serum starvation and       |
|                     | etoposide-induced apoptosis. The mechanism through which TDE1 protects cells is poorly        |
|                     | understood, but may involve aberrant methylation of TDE1 complexes.                           |
| Gene ID:            | 10955   |
| Application Details |   |
| Application Notes:  | IF(IHC-P) 1:50-200  |
|                     | IF(IHC-F) 1:50-200  |
|                     | IF(ICC) 1:50-200  |
| Restrictions:       | For Research Use only   |
| Handling            |   |
| Format:             | Liquid  |
| Concentration:      | 1 μg/μL   |
| Buffer:             | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and   |
|                     | 50 % Glycerol.  |
| Preservative:       | ProClin   |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be           |
|                     | handled by trained staff only.  |
| Storage:            | -20 °C  |
| Storage Comment:    | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.             |
| Expiry Date:        | 12 months   |
|                     |   |