

Datasheet for ABIN7601680
anti-ALDH3A1 antibody (AA 413-453)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	ALDH3A1
Binding Specificity:	AA 413-453
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ALDH3A1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Purpose:	Anti-ALDH3A1 Antibody Picoband®
Immunogen:	E.coli-derived human ALDH3A1 recombinant protein (Position: Y413-H453).
Isotype:	IgG
Cross-Reactivity (Details):	No cross-reactivity with other proteins.
Characteristics:	Anti-ALDH3A1 Antibody Picoband® (ABIN7601680). Tested in ELISA, IF, IHC, ICC, WB applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Purification:	Immunogen affinity purified.

Target Details

Target:	ALDH3A1
Alternative Name:	ALDH3A1 (ALDH3A1 Products)
Background:	Synonyms: CREB-regulated transcription coactivator 2, Transducer of regulated cAMP response element-binding protein 2, TORC-2, Transducer of CREB protein 2, CRTC2, TORC2 Tissue Specificity: Most abundantly expressed in the thymus. Present in both B and T-lymphocytes. Highly expressed in HEK293T cells and in insulinomas. High levels also in spleen, ovary, muscle and lung, with highest levels in muscle. Lower levels found in brain, colon, heart, kidney, prostate, small intestine and stomach. Weak expression in liver and pancreas. Background: ALDH3A1(Aldehyde Dehydrogenase, Family 3, Subfamily A, Member 1), also known as ALDH3, is an enzyme that in humans is encoded by the ALDH3A1 gene. By in situ hybridization, Hiraoka et al.(1995) mapped the ALDH3 gene to 17p11.2. ALDHs play a major role in the detoxification of alcohol-derived acetaldehyde. They are involved in the metabolism of corticosteroids, biogenic amines, neurotransmitters, and lipid peroxidation. This protein preferentially oxidizes aromatic aldehyde substrates. It may play a role in the oxidation of toxic aldehydes.
Molecular Weight:	55 kDa
Gene ID:	218
UniProt:	P30838

Application Details

Application Notes:	Western blot, 0.25-0.5 µg/mL, Human Immunohistochemistry(Paraffin-embedded Section), 2-5 µg/mL, Human Immunocytochemistry/Immunofluorescence, 5 µg/mL, Human ELISA, 0.1-0.5 µg/mL, - 1. Hiraoka, L. R., Hsu, L., Hsieh, C.-L. Assignment of ALDH3 to human chromosome 17p11.2 and ALDH5 to human chromosome 9p13. Genomics 25: 323-325, 1995. 2. Santisteban, I., Povey, S., West, L. F., Parrington, J. M., Hopkinson, D. A. Chromosome assignment, biochemical and immunological studies on a human aldehyde dehydrogenase, ALDH3. Ann. Hum. Genet. 49: 87-100, 1985.
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
---------	-------------

Handling

Reconstitution:	Adding 0.2 mL of distilled water will yield a concentration of 500 µg/mL.
Concentration:	500 µg/mL
Buffer:	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage:	4 °C, -20 °C
Storage Comment:	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.