

Datasheet for ABIN7601738
anti-PNP antibody (AA 44-273)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	PNP
Binding Specificity:	AA 44-273
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PNP antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Flow Cytometry (FACS)

Product Details

Purpose:	Anti-PNP Antibody Picoband®
Immunogen:	E.coli-derived human PNP recombinant protein (Position: Q44-Q273). Human PNP shares 86.5% and 87% amino acid (aa) sequence identity with mouse and rat PNP, respectively.
Characteristics:	Anti-PNP Antibody Picoband® (ABIN7601738). Tested in WB, Flow Cytometry, ELISA applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Purification:	Immunogen affinity purified.

Target Details

Target:	PNP
Alternative Name:	PNP (PNP Products)
Background:	The PNP gene encodes purine nucleoside phosphorylase, an enzyme that catalyzes the reversible phosphorolysis of the purine nucleosides and deoxynucleosides inosine, guanosine, deoxyinosine, and deoxyguanosine. It is presented results from gene dosage studies consistent with assignment of the PNP locus to band 14q13. PNP is expressed in most tissues, with markedly greater expression in lymphoid tissues. Genetic deficiencies of PNP result in severely compromised T γ lymphocyte function and neurologic dysfunction.
Molecular Weight:	32 kDa
Gene ID:	4860
UniProt:	P00491
Pathways:	Regulation of Leukocyte Mediated Immunity , Positive Regulation of Immune Effector Process , Ribonucleoside Biosynthetic Process , Positive Regulation of Response to DNA Damage Stimulus

Application Details

Application Notes:	Western blot, 0.1-0.25 μ g/mL, Human, Mouse, Rat Flow Cytometry (Fixed), 1-3 μ g/1 $\times$ 10 ⁶ cells, Human ELISA, 0.1-0.5 μ g/mL, - 1. Williams, S. R., Goddard, J. M., Martin, D. W., Jr. Human purine nucleoside phosphorylase cDNA sequence and genomic clone characterization. Nucleic Acids Res. 12: 5779-5787, 1984. 2. Frecker, M., Dallaire, L., Young, S. R., Chen, G. C. C., Simpson, N. E. Confirmation of regional assignment of nucleoside phosphorylase (NP) on chromosome 14 by gene dosage studies. Hum. Genet. 45: 167-173, 1978. 3. Markert, M. L., Finkel, B. D., McLaughlin, T. M., Watson, T. J., Collard, H. R., McMahon, C. P., Andrews, L. G., Barrett, M. J., Ward, F. E. Mutations in purine nucleoside phosphorylase deficiency. Hum. Mutat. 9: 118-121, 1997.
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Adding 0.2 mL of distilled water will yield a concentration of 500 μ g/mL.
Concentration:	500 μ g/mL

Handling

Buffer:	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage:	4 °C,-20 °C
Storage Comment:	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.