

Datasheet for ABIN7601796
anti-HYI antibody (AA 46-277)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	HYI
Binding Specificity:	AA 46-277
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HYI antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Flow Cytometry (FACS), Immunocytochemistry (ICC), Immunofluorescence (IF)

Product Details

Purpose:	Anti-HYI Antibody Picoband®
Immunogen:	E.coli-derived human HYI recombinant protein (Position: E46-Q277).
Isotype:	IgG
Cross-Reactivity (Details):	No cross-reactivity with other proteins
Characteristics:	Anti-HYI Antibody Picoband® (ABIN7601796). Tested in ELISA, Flow Cytometry, IF, IHC, ICC, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Purification:	Immunogen affinity purified.

Target Details

Target:	HYI
Alternative Name:	HYI (HYI Products)
Background:	Synonyms: RNA-binding protein 47,RNA-binding motif protein 47,RBM47, Tissue Specificity: Abundantly expressed in tonsil, lymph node, and trachea, strong expression in prostate, lower expression in thyroid, stomach, and colon. . Background: This gene encodes a putative hydroxypyruvate isomerase, which likely catalyzes the conversion of hydroxypyruvate to 2-hydroxy-3-oxopropanoate, and may be involved in carbohydrate transport and metabolism. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.
Molecular Weight:	30 kDa
Gene ID:	81888
UniProt:	Q5T013

Application Details

Application Notes:	Western blot, 0.25-0.5 µg/mL, Mouse, Rat Immunohistochemistry (Paraffin-embedded Section), 2-5 µg/mL, Human Immunocytochemistry/Immunofluorescence, 5 µg/mL, Human Flow Cytometry (Fixed), 1-3 µg/1x10⁶ cells, Human ELISA, 0.1-0.5 µg/mL, - 1. Arunachalam, B., Phan, U. T., Geuze, H. J., Cresswell, P. Enzymatic reduction of disulfide bonds in lysosomes: characterization of a gamma-interferon-inducible lysosomal thiol reductase (GILT). Proc. Nat. Acad. Sci. 97: 745-750, 2000. 2. Maric, M., Arunachalam, B., Phan, U. T., Dong, C., Garrett, W. S., Cannon, K. S., Alfonso, C., Karlsson, L., Flavell, R. A., Cressell, P. Defective antigen processing in GILT-free mice. Science 294: 1361-1365, 2001. 3. Phan, U. T., Arunachalam, B., Cresswell, P. Gamma-interferon-inducible lysosomal thiol reductase (GILT): maturation, activity, and mechanism of action. J. Biol. Chem. 275: 25907-25914, 2000.
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Add 0.2 mL of distilled water will yield a concentration of 500 µg/mL.
Concentration:	500 µg/mL

Handling

Buffer:	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ , 0.01 mg Sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.