

Datasheet for ABIN7602149
anti-TXN2 antibody (AA 60-166)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	TXN2
Binding Specificity:	AA 60-166
Reactivity:	Human, Rat, Mouse
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This TXN2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC), Flow Cytometry (FACS)

Product Details

Purpose:	Anti-Thioredoxin 2/TXN2 Antibody Picoband® (monoclonal, 7B5)
Immunogen:	E.coli-derived human Thioredoxin 2/TXN2 recombinant protein (Position: T60-G166).
Clone:	7B5
Isotype:	IgG2b
Cross-Reactivity (Details):	No cross-reactivity with other proteins.
Characteristics:	Anti-Thioredoxin 2/TXN2 Antibody Picoband® (monoclonal, 7B5) (ABIN7602149). Tested in Flow Cytometry, IF, IHC, ICC, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.

Product Details

Purification: Immunogen affinity purified.

Target Details

Target: TXN2

Alternative Name: TXN2 ([TXN2 Products](#))

Background: Synonyms: Thioredoxin, mitochondrial, MTRX, Mt-Trx, Thioredoxin-2, TXN2, TRX2
Tissue Specificity: Widely expressed in adult (at protein level) and fetal tissues.
Background: Thioredoxin, mitochondrial also known as thioredoxin-2 is a protein that in humans is encoded by the TXN2 gene on chromosome 22. It is mapped to 22q12.3. This nuclear gene encodes a mitochondrial member of the thioredoxin family, a group of small multifunctional redox-active proteins. The encoded protein may play important roles in the regulation of the mitochondrial membrane potential and in protection against oxidant-induced apoptosis.

Molecular Weight: 14 kDa

Gene ID: 25828

UniProt: [Q99757](#)

Pathways: [Cell RedoxHomeostasis](#)

Application Details

Application Notes: Western blot, 0.1-0.5 µg/mL, Human, Mouse, Rat
Immunohistochemistry (Paraffin-embedded Section), 0.5-1 µg/mL, Human, Mouse, Rat
Immunocytochemistry/Immunofluorescence, 2 µg/mL, Human
Flow Cytometry (Fixed), 1-3 µg/1x10⁶ cells, Human
1. Benhar, M., Forrester, M. T., Hess, D. T., Stamler, J. S. Regulated protein denitrosylation by cytosolic and mitochondrial thioredoxins. Science 320: 1050-1054, 2008. 2. Chen, Y., Cai, J., Murphy, T. J., Jones, D. P. Overexpressed human mitochondrial thioredoxin confers resistance to oxidant-induced apoptosis in human osteosarcoma cells. J. Biol. Chem. 277: 33242-33248, 2002.

Restrictions: For Research Use only

Handling

Format: Lyophilized

Handling

Reconstitution:	Add 0.2 mL of distilled water will yield a concentration of 500 µg/mL.
Concentration:	500 µg/mL
Buffer:	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ , 0.05 mg NaN ₃ .
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.