

Datasheet for ABIN7602578
anti-ZNF609 antibody (AA 865-1283)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	ZNF609
Binding Specificity:	AA 865-1283
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZNF609 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Flow Cytometry (FACS), Immunocytochemistry (ICC), Immunofluorescence (IF)

Product Details

Purpose:	Anti-ZNF609 Antibody Picoband®
Immunogen:	E.coli-derived human ZNF609 recombinant protein (Position: K865-R1283).
Isotype:	IgG
Cross-Reactivity (Details):	No cross-reactivity with other proteins.
Characteristics:	Anti-Z Antibody Picoband® (ABIN7602578). Tested in ELISA, Flow Cytometry, IF, ICC, WB applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Purification:	Immunogen affinity purified.

Target Details

Target:	ZNF609
Alternative Name:	ZNF609 (ZNF609 Products)
Background:	<p>Synonyms: Kelch repeat and BTB domain-containing protein 2, BTB and kelch domain-containing protein 1, KBTBD2, BKLHD1, KIAA1489</p> <p>Tissue Specificity: Detected in liver, skeletal muscle, kidney, pancreas, spleen, thyroid, testis, ovary, small intestine and colon.</p> <p>Background: Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. zinc finger protein 609 (ZNF609) is a 1,411 amino acid member of the Krüppel C2H2-type zinc-finger protein family. Localized to the nucleus, ZNF609 contains one C2H2-type zinc finger through which it is thought to be involved in DNA-binding and transcriptional regulation.</p>
Molecular Weight:	180 kDa
Gene ID:	23060
UniProt:	O15014

Application Details

Application Notes:	<p>Western blot, 0.25-0.5 µg/mL, Human</p> <p>Immunocytochemistry/Immunofluorescence, 5 µg/mL, Human</p> <p>Flow Cytometry (Fixed), 1-3 µg/1x10⁶ cells, Human</p> <p>ELISA, 0.1-0.5 µg/mL, -</p> <p>1. Legnini, I., Di Timoteo, G., Rossi, F., Morlando, M., Briganti, F., Sthandier, O., Fatica, A., Santini, T., Andronache, A., Wade, M., Laneve, P., Rajewsky, N., Bozzoni, I. Circ-ZNF609 is a circular RNA that can be translated and functions in myogenesis. Molec. Cell 66: 22-37, 2017. 2. Nagase, T., Ishikawa, K., Nakajima, D., Ohira, M., Seki, N., Miyajima, N., Tanaka, A., Kotani, H., Nomura, N., Ohara, O. Prediction of the coding sequences of unidentified human genes. VII. The complete sequences of 100 new cDNA clones from brain which can code for large proteins in vitro. DNA Res. 4: 141-150, 1997.</p>
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
---------	-------------

Handling

Reconstitution:	Adding 0.2 mL of distilled water will yield a concentration of 500 µg/mL.
Concentration:	500 µg/mL
Buffer:	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage:	4 °C, -20 °C
Storage Comment:	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.