

Datasheet for ABIN7602637
anti-DBN1 antibody (AA 9-649)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	DBN1
Binding Specificity:	AA 9-649
Reactivity:	Human, Mouse, Rat, Monkey
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This DBN1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Flow Cytometry (FACS)

Product Details

Purpose:	Anti-Drebrin/DBN1 Antibody Picoband® (monoclonal, 4F6E7)
Immunogen:	E.coli-derived human Drebrin/DBN1 recombinant protein (Position: H9-D649).
Clone:	4F6E7
Isotype:	IgG2a
Cross-Reactivity (Details):	No cross-reactivity with other proteins.
Characteristics:	Anti-Drebrin/DBN1 Antibody Picoband® (monoclonal, 4F6E7) (ABIN7602637). Tested in Flow Cytometry, IHC, WB applications. This antibody reacts with Human, Mouse, Rat, Monkey. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.

Product Details

Purification: Immunogen affinity purified.

Target Details

Target: DBN1

Alternative Name: DBN1 ([DBN1 Products](#))

Background: Synonyms: Cytochrome c oxidase subunit 4 isoform 1, mitochondrial, Cytochrome c oxidase polypeptide IV, Cytochrome c oxidase subunit IV isoform 1, COX IV-1, COX4I1, COX4
Tissue Specificity: Highly expressed in liver. Expressed in testis and weakly in colon.
Background: Drebrin is a protein that in humans is encoded by the DBN1 gene. The protein encoded by this gene is a cytoplasmic actin-binding protein thought to play a role in the process of neuronal growth. It is a member of the drebrin family of proteins that are developmentally regulated in the brain. A decrease in the amount of this protein in the brain has been implicated as a possible contributing factor in the pathogenesis of memory disturbance in Alzheimer's disease. At least two alternative splice variants encoding different protein isoforms have been described for this gene.

Molecular Weight: 120 kDa

Gene ID: 1627

UniProt: [Q16643](#)

Pathways: [Maintenance of Protein Location](#)

Application Details

Application Notes: Western blot, 0.25-0.5 µg/mL, Human, Mouse, Rat, Monkey
Immunohistochemistry(Paraffin-embedded Section), 2-5 µg/mL, Human, Rat
Flow Cytometry (Fixed), 1-3 µg/1x10⁶ cells, Human
1. Akbar, M., Kim, H. Y. Protective effects of docosahexaenoic acid in staurosporine-induced apoptosis: involvement of phosphatidylinositol-3 kinase pathway. J. Neurochem. 82: 655-665,
2. Calon, F., Lim, G. P., Yang, F., Morihara, T., Teter, B., Ubeda, O., Rostaing, P., Triller, A., Salem, N., Jr., Ashe, K. H., Frautschy, S. A., Cole, G. M. Docosahexaenoic acid protects from dendritic pathology in an Alzheimer's disease mouse model. Neuron 43: 633-645, 2004. 3. Harigaya, Y., Shoji, M., Shirao, T., Hirai, S. Disappearance of actin-binding protein, drebrin, from hippocampal synapses in Alzheimer's disease. J. Neurosci. Res. 43: 87-92, 1996.

Restrictions: For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Adding 0.2 mL of distilled water will yield a concentration of 500 µg/mL.
Concentration:	500 µg/mL
Buffer:	Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na ₂ HPO ₄ .
Storage:	4 °C, -20 °C
Storage Comment:	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.