

Datasheet for ABIN7602916 **anti-RPL32 antibody (C-Term)**



[Go to Product page](#)

Overview

| | |
|----------------------|--|
| Quantity: | 100 µg |
| Target: | RPL32 |
| Binding Specificity: | C-Term |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This RPL32 antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunofluorescence (IF), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Flow Cytometry (FACS) |

Product Details

| | |
|-----------------------------|--|
| Purpose: | Anti-RPL32 Antibody Picoband® |
| Immunogen: | A synthetic peptide corresponding to a sequence at the C-terminus of human RPL32, identical to the related mouse and rat sequences. |
| Isotype: | IgG |
| Cross-Reactivity (Details): | No cross-reactivity with other proteins. |
| Characteristics: | Anti-RPL32 Antibody Picoband® (ABIN7602916). Tested in Flow Cytometry, IF, IHC, ICC, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance. |

Product Details

Purification: Immunogen affinity purified.

Target Details

Target: RPL32

Alternative Name: RPL32 ([RPL32 Products](#))

Background: Synonyms: Max dimerization protein 1, Max dimerizer 1, Protein MAD, MXD1, MAD
Background: 60S ribosomal protein L32 is a protein that in humans is encoded by the RPL32 gene. Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein belongs to the L32E family of ribosomal proteins. It is located in the cytoplasm. Although some studies have mapped this gene to 3q13.3-q21, it is believed to map to 3p25-p24. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome. Alternatively spliced transcript variants encoding the same protein have been observed for this gene.

Molecular Weight: 18 kDa

Gene ID: 6161

UniProt: [P62910](#)

Application Details

Application Notes: Western blot, 0.1-0.25 µg/mL/mL, Human, Mouse, Rat
Immunohistochemistry(Paraffin-embedded Section), 2-5 µg/mL/mL, Human
Immunocytochemistry/Immunofluorescence, 5 µg/mL/mL, Human
Immunofluorescence, 5 µg/mL/mL, Human
Flow Cytometry (Fixed), 1-3 µg/mL/1x10⁶ cells, Human
1. Neto, E. D. , Correa, R. G. , & Verjovski-Almeida, S. . (2000). Shotgun sequencing of the human transcriptome with orf expressed sequence tags. Proceedings of the National Academy of Sciences, 97(7), 3491-3496.

Restrictions: For Research Use only

Handling

Format: Lyophilized

Handling

| | |
|------------------|--|
| Reconstitution: | Adding 0.2 mL of distilled water will yield a concentration of 500 µg/mL. |
| Concentration: | 500 µg/mL |
| Buffer: | Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ . |
| Storage: | 4 °C, -20 °C |
| Storage Comment: | At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing. |