

Datasheet for ABIN7603106
anti-MT-ND2 antibody (N-Term)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	MT-ND2
Binding Specificity:	N-Term
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MT-ND2 antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS)

Product Details

Purpose:	Anti-NADH2/Mtnd2 Antibody Picoband®
Immunogen:	A synthetic peptide corresponding to a sequence at the N-terminus of rat NADH2/Mtnd2, which shares 80% amino acid (aa) sequence identity with mouse NADH2/Mtnd2.
Isotype:	IgG
Cross-Reactivity (Details):	No cross-reactivity with other proteins.
Characteristics:	Anti-NADH2/Mtnd2 Antibody Picoband® (ABIN7603106). Tested in Flow Cytometry, WB applications. This antibody reacts with Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Purification:	Immunogen affinity purified.

Target Details

Target:	MT-ND2
Alternative Name:	Mtnd2 (MT-ND2 Products)
Background:	<p>Synonyms: NADH-ubiquinone oxidoreductase chain 2, NADH dehydrogenase subunit 2, Mtnd2, mt-Nd2, Nd2</p> <p>Tissue Specificity: Ubiquitously expressed.</p> <p>Background: Mitochondrially encoded NADH dehydrogenase 2 is protein that in humans is encoded by the mitochondrial gene MT-ND2 gene. The ND2 protein is a subunit of NADH dehydrogenase (ubiquinone), which is located in the mitochondrial inner membrane and is the largest of the five complexes of the electron transport chain. Variants of MT-ND2 are associated with mitochondrial encephalomyopathy, lactic acidosis, and stroke-like episodes (MELAS), Leigh's syndrome (LS), Leber's hereditary optic neuropathy (LHON) and increases in adult BMI.</p>
Molecular Weight:	39 kDa
Gene ID:	26194
UniProt:	P11662

Application Details

Application Notes:	<p>Western blot, 0.25-0.5 µg/mL, Mouse, Rat</p> <p>Flow Cytometry (Fixed), 1-3 µg/1×10⁶ cells, Rat</p> <p>1. Anderson, S., Bankier, A. T., Barrell, B. G., de Bruijn, M. H. L., Coulson, A. R., Drouin, J., Eperon, I. C., Nierlich, D. P., Roe, B. A., Sanger, F., Schreier, P. H., Smith, A. J. H., Staden, R., Young, I. G. Sequence and organization of the human mitochondrial genome. Nature 290: 457-465, 1981. 2. Arizmendi, J. M., Skehel, J. M., Runswick, M. J., Fearnley, I. M., Walker, J. E. Complementary DNA sequences of two 14.5 kDa subunits of NADH:ubiquinone oxidoreductase from bovine heart mitochondria. Complementation of the primary structure of the complex FEBS Lett. 313: 80-84, 1992. 3. Attardi, G., Chomyn, A., Doolittle, R. F., Mariottini, P., Ragan, C. I. Seven unidentified reading frames of human mitochondrial DNA encode subunits of the respiratory chain NADH dehydrogenase. Cold Spring Harbor Symp. Quant. Biol. 1: 103-114, 1986.</p>
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
---------	-------------

Handling

Reconstitution:	Add 0.2 mL of distilled water will yield a concentration of 500 µg/mL.
Concentration:	500 µg/mL
Buffer:	Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na ₂ HPO ₄ .
Storage:	4 °C, -20 °C
Storage Comment:	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.