-online.com antibodies

Datasheet for ABIN760436 anti-CDCA7L antibody (AA 381-454)



Overview

Quantity:	100 μL	
Target:	CDCA7L	
Binding Specificity:	AA 381-454	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CDCA7L antibody is un-conjugated	
Application:	ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin- embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CDCA7L	
Isotype:	lgG	
Cross-Reactivity:	Human, Mouse, Rat	
Predicted Reactivity:	Dog,Cow,Pig,Horse,Chicken,Rabbit	
Purification:	Purified by Protein A.	
Target Details		
Target:	CDCA7L	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN760436 | 03/06/2024 | Copyright antibodies-online. All rights reserved.

Background: Synonyms: c Myc Interacting protein JPO2, DKFZp762L0311, RAM2, HR1, JPO2, Protein JPO2, R1 antibody Transcription Factor RAM2 CDA7L_HUMAN Background: CDCA7L. Is a member of the CDCA (cell division cycle associated) protein family. Members of this family have expression patterns associated with other known cell cycle genes such as cyclins and CDC genes. CDCA7L plays a role in transcriptional regulation as a represso that inhibits monoamine oxidase A (MAOA) activity and gene expression by binding to the promoter. It also plays an important oncogenic role in mediating the full transforming effect of MYC in medialbalstoma cells. It is involved in apoptotic signaling pathways and may act downstream of P38-kinase and BCL-2, but upstream of CASP3/caspase-3 as well as CCND1/cyclin D1 and E2F1. Gene ID: S5536 Application Details ELISA 1.500-1000 HiCF 11:00-500 IFC(HC-F) 1:50-200 IF(HC -F) 1:50-200 IF(HC -F) 1:50-200 IF(HC -F) 1:50-200 IFC(HC -F) 1:50-200 IFG(HC -F) 1:50-200 IFC(HC -F) 1:50-200 IFG(HC -F) 1:50-200 IFC(HC -F) 1:50-200 Bardititions: For Research Use on	Target Details		
R1 antibody Transcription Factor RAM2 CDA7L HUMAN Background: CDCA7L is a member of the CDCA (cell division cycle associated) protein family. Members of this family have expression patterns associated with other known cell cycle genes such as cyclins and CDC genes. CDCA7L plays a role in transcriptional regulation as a represso that inhibits monoamine oxidase A (MAOA) activity and gene expression by binding to the promoter. It also plays an important oncogenic role in mediating the full transforming effect of MYC in medulloblastoma cells. It is involved in apoptotic signaling pathways and may act downstream of P38-kinase and BCL-2, but upstream of CASP3/caspase-3 as well as CCND1/cyclin D1 and E2F1. Gene ID: 55536 Application Details ELISA 1:500-1000 HC-F1 1200-400 HC-F1 1:00-500 IF(HC-P) 1:50-200 IF(HC-P) 1:50-200 IF(HC-F) 1:50-200 IF((HC-P) 1:50-200 IF((CC) 1:50-200 IF((CC) 1:50-200 Buffer: 0.01M TBS(pH7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	Alternative Name:	Cdca7l (CDCA7L Products)	
Background: CDCA7L is a member of the CDCA (cell division cycle associated) protein family. Members of this family have expression patterns associated with other known cell cycle genes such as cyclins and CDC genes. CDCA7L plays a role in transcriptional regulation as a repression that inhibits monoamine oxidase A (MAOA) activity and gene expression by binding to the promoter. It also plays an important oncogenic role in mediating the full transforming effect of MYC in medulioblastoma cells. It is involved in apoptotic signaling pathways and may act downstream of P38-kinase and BCL-2, but upstream of CASP3/caspase-3 as well as CCND1/cyclin D1 and E2F1. Gene ID: 55536 Application Details ELISA 1.500-1000 IHC-P 1.200-400 IHC-P 1.50-200 IF(HC-P) 1.50-200 IF(HC-P) 1.50-200 IF(HC-P) 1.50-200 IF(HC-P) 1.50-200 IF(ICC) 1.50-200 IF(ICC) 1.50-200 Restrictions: For Research Use only Handling 1 µg/µL Buffer: 0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % ProcIn300 and 50 % Glycerol. Preservative: ProClin Preservative: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	Background:	Synonyms: c Myc Interacting protein JPO2, DKFZp762L0311, RAM2, HR1, JPO2, Protein JPO2,	
Members of this family have expression patterns associated with other known cell cycle genes such as cyclins and CDC genes. CDCA7L plays a role in transcriptional regulation as a repression that inhibits monoamine oxidase A (MAOA) activity and gene expression by binding to the promoter. It also plays an important oncogenic role in mediating the full transforming effect of MYC in medulloblastoma cells. It is involved in apoptotic signaling pathways and may act downstream of P38-kinase and BCL-2, but upstream of CASP3/caspase-3 as well as CCND1/cyclin D1 and E2F1. Gene ID: 55536 Application Details ELISA 1:500-1000 IHC-F 1:200-400 IHC-F 1:200-400 IF(HC-F) 1:50-200 IF(HC-F) 1:50-200		R1 antibody Transcription Factor RAM2.CDA7L_HUMAN	
such as cyclins and CDC genes. CDCA7L plays a role in transcriptional regulation as a represso that inhibits monoamine oxidase A (MAOA) activity and gene expression by binding to the promoter. It also plays an important oncogenic role in mediating the full transforming effect of MYC in medulloblastoma cells. It is involved in apoptotic signaling pathways and may act downstream of P38-kinase and BCL-2, but upstream of DASP3/caspase-3 as well as COND1/cyclin D1 and E2F1. Gene ID: \$5536 Application Details Application Notes: ELISA 1:500-1000 IHC-F 1:100-500 IF(HC-F) 1:50-200 AndIting Format: Liquid Concentration: 1 µg/µL Buffer: 0.011M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol. Preservative: ProClin Proclation of Use: This product contains ProClin: a POISONOUS AND HA		Background: CDCA7L is a member of the CDCA (cell division cycle associated) protein family.	
that inhibits monoamine oxidase A (MAOA) activity and gene expression by binding to the promoter. It also plays an important oncogenic role in mediating the full transforming effect of MYC in meduiloblastoma cells. It is involved in apoptotic signaling pathways and may act downstream of P38-kinase and BCL-2, but upstream of CASP3/caspase-3 as well as CCND1/cyclin D1 and E2F1.Gene ID:55536Application DetailsApplication Notes:ELISA 1:500-1000 IHC-P 1:200-400 IHC-P 1:200-400 IF(IHC-P) 1:50-200 IF(IHC-P) 1:50-200 IF(IHC-P) 1:50-200 IF(ICC) 1:50-200Restrictions:For Research Use onlyHandlingFormat:LiquidConcentration:1 µg/µLBuffer:0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.Precaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.		Members of this family have expression patterns associated with other known cell cycle genes	
promoter. It also plays an important oncogenic role in mediating the full transforming effect of MYC in medulloblastoma cells. It is involved in apoptotic signaling pathways and may act downstream of P38-kinase and BCL-2, but upstream of CASP3/caspase-3 as well as CCND1/cyclin D1 and E2F1. Gene ID: 55536 Application Details ELISA 1:500-1000 IHC-P 1:200-400 IHC-P 1:200-400 IFC(HC-P) 1:50-200 IFC(HC-P) 1:50-200 		such as cyclins and CDC genes. CDCA7L plays a role in transcriptional regulation as a repressor	
MYC in medulioblastoma cells. It is involved in apoptotic signaling pathways and may act downstream of P38-kinase and BCL-2, but upstream of CASP3/caspase-3 as well as CCND1/cyclin D1 and E2F1. Gene ID: 55536 Application Details Application Notes: ELISA 1:500-1000 IHC-P 1:200-400 IHC-P 1:200-400 IHC-F) 1:50-200 IF((HC-F) 1:50-200 IF((HC-F) 1:50-200 IF((CC) 1:50-200 IF((CC) 1:50-200 IF((CC) 1:50-200 IF(ICC) 1:50-200 IF(ICC) 1:50-200 Handling IF(ICC) 1:50-200 Promat: Liquid Concentration: 1 µg/µL Buffer: 0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol. Preservative: ProClin Preceution of		that inhibits monoamine oxidase A (MAOA) activity and gene expression by binding to the	
downstream of P38-kinase and BCL-2, but upstream of CASP3/caspase-3 as well as CCND1/cyclin D1 and E2F1. Gene ID: 55536 Application Details Application Notes: ELISA 1:500-1000 IHC-F 1:200-400 IHC-F 1:50-200 IF(IHC-P) 1:50-200 IF(IHC-P) 1:50-200 IF(ICC) 1:50-200 IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: 0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol. Preservative: ProClin Preservative: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. Storage: 4 *0, 20 *0			
CCND1/cyclin D1 and E2F1. Gene ID: 55536 Application Details Application Notes: ELISA 1:500-1000 IHC-P 1:200-400 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-P) 1:50-200 IF(IHC-P) 1:50-200 Restrictions: For Research Use only Handling Iuquid Format: Liquid Concentration: 1 µg/µL Buffer: 0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol. Preservative: ProClin Preservative: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. Storage: 4 "C-20 "C			
Gene ID: 55536 Application Details ELISA 1:500-1000 Application Notes: ELISA 1:500-1000 IHC-P 1:200-400 IHC-P 1:200-400 IHC-F 1:100-500 IF((IHC-P) 1:50-200 IF((IHC-P) 1:50-200 IF((IHC-P) 1:50-200 IF((IHC-P) 1:50-200 IF((IHC-P) 1:50-200 IF(IHC-P) 1:50-200 IF(IHC-P) 1:50-200 IF(IHC-P) 1:50-200 IF(IHC-P) 1:50-200 IF(IHC-P) 1:50-200 IF(IHC-P) 1:50-200 Restrictions: For Research Use only Handling Iuquid Concentration: Iuquid Concentration: 1 µg/µL Buffer: 0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. Storage: 4'C, 20 'C			
Application DetailsApplication Notes:ELISA 1:500-1000 HG-P 1:200-400 HG-F 1:100-500 IF(HC-P) 1:50-200 IF(HC-P) 1:50-200 IF(ICC) 1:50-200Restrictions:For Research Use onlyHandlingFormat:LiquidConcentration:1 µg/µLBuffer:0.11M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.Preservative:ProClinPreservative:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.Storage:4 °C-20 °C		CCND1/cyclin D1 and E2F1.	
Application Notes:ELISA 1:500-1000 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-F) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200Restrictions:For Research Use onlyHandlingIiquidFormat:LiquidConcentration:1 µg/µLBuffer:0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.Preservative:ProClinPrecaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.Storage:4 °C-20 °C	Gene ID:	55536	
IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-F) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200Restrictions:For Research Use onlyHandlingIuquidFormat:LiquidConcentration:1 µg/µLBuffer:0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.Preservative:ProClinPreservative:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.Storage:4 °C-20 °C	Application Details		
HC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200Restrictions:For Research Use onlyHandlingIuquidFormat:LiquidConcentration:1 μg/μLBuffer:0.11M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.Preservative:ProClinPreservative:No ClinPreservative:Ais product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.Storage:4°.20 °C	Application Notes:	ELISA 1:500-1000	
IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200Restrictions:For Research Use onlyHandlingIuquidFormat:LiquidConcentration:1 μg/μLBuffer:0.11M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.Preservative:ProClinPrecaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.Storage:4 °C-20 °C		IHC-P 1:200-400	
IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 μg/μL Buffer: 0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. Storage: 4 °C,-20 °C		IHC-F 1:100-500	
IF(ICC) 1:50-200Restrictions:For Research Use onlyHandlingFormat:LiquidConcentration:1 μg/μLBuffer:0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.Preservative:ProClinPrecaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.Storage:4 °C, 20 °C		IF(IHC-P) 1:50-200	
Restrictions:For Research Use onlyHandlingFormat:LiquidConcentration:1 µg/µLBuffer:0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.Preservative:ProClinPrecaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.Storage:4 °C, 20 °C		IF(IHC-F) 1:50-200	
HandlingFormat:LiquidConcentration:1 μg/μLBuffer:0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.Preservative:ProClinPrecaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.Storage:4 °C,-20 °C		IF(ICC) 1:50-200	
Format:LiquidConcentration:1 μg/μLBuffer:0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.Preservative:ProClinPrecaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.Storage:4 °C,-20 °C	Restrictions:	For Research Use only	
Concentration:1 µg/µLBuffer:0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.Preservative:ProClinPrecaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.Storage:4 °C,-20 °C	Handling		
Buffer:0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.Preservative:ProClinPrecaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.Storage:4 °C,-20 °C	Format:	Liquid	
Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. Storage: 4 °C,-20 °C	Concentration:	1 μg/μL	
Precaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.Storage:4 °C,-20 °C	Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
handled by trained staff only. Storage: 4 °C,-20 °C	Preservative:	ProClin	
Storage: 4 °C,-20 °C	Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
		handled by trained staff only.	
Storage Comment: Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.	Storage:	4 °C,-20 °C	
	Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN760436 | 03/06/2024 | Copyright antibodies-online. All rights reserved.

1.1	(1:
Н	land	ling
		3

Expiry Date:

12 months

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN760436 | 03/06/2024 | Copyright antibodies-online. All rights reserved.