

## Datasheet for ABIN7605208

## anti-DFFA antibody



| ()      | ve  | r\/i | Δ | ۱۸/ |
|---------|-----|------|---|-----|
| $\circ$ | V C | 1 V  |   | v v |

| Quantity:    | 100 μL   |  |
|--------------|--|--|
| Target:      | DFFA   |  |
| Reactivity:  | Human  |  |
| Host:        | Rabbit   |  |
| Clonality:   | Monoclonal   |  |
| Conjugate:   | This DFFA antibody is un-conjugated  |  |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunoprecipitation (IP), Immunocytochemistry (ICC), Flow Cytometry (FACS) |  |

## **Product Details**

| Purpose:         | Anti-ICAD Rabbit Monoclonal Antibody   |  |
|------------------|--|--|
| Immunogen:       | A synthesized peptide derived from human ICAD  |  |
| Clone:           | 21D96  |  |
| Isotype:         | IgG  |  |
| Characteristics: | Anti-ICAD Rabbit Monoclonal Antibody (ABIN7605208). Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human. |  |
| Purification:    | Affinity-chromatography  |  |
| Target Dataila   |  |  |

Target Details

Target: DFFA

## **Target Details**

| rarget Details      |  |  |
|---------------------|--|--|
| Alternative Name:   | DFFA (DFFA Products)   |  |
| Background:         | Synonyms: Cubilin,460 kDa receptor,Intestinal intrinsic factor receptor,Intrinsic factor-        |  |
|                     | cobalamin receptor,Intrinsic factor-vitamin B12 receptor,CUBN,IFCR,                              |  |
|                     | Tissue Specificity: Expressed in kidney proximal tubule cells, placenta, visceral yolk-sac cells |  |
|                     | and in absorptive intestinal cells. Expressed in the epithelium of intestine and kidney.         |  |
| Molecular Weight:   | 37 kDa   |  |
| UniProt:            | 000273   |  |
| Pathways:           | Apoptosis, Caspase Cascade in Apoptosis  |  |
| Application Details |  |  |
| Application Notes:  | WB 1:500-1:2000  |  |
|                     | IHC 1:50-1:200   |  |
|                     | ICC/IF 1:50-1:200  |  |
|                     | IP 1:50  |  |
|                     | FC 1:50  |  |
| Restrictions:       | For Research Use only  |  |
| Handling            |  |  |
| Format:             | Liquid   |  |
| Reconstitution:     | Restore with deionized water (or equivalent) for reconstitution volume of 1.0 mL                 |  |
| Concentration:      | Lot specific   |  |
| Buffer:             | Rabbit IgG in phosphate buffered saline, pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 %       |  |
|                     | glycerol, 0.4-0.5 mg/mL BSA.   |  |
| Preservative:       | Sodium azide   |  |
| Precaution of Use:  | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which                    |  |
|                     | should be handled by trained staff only.   |  |
| Storage:            | 4 °C,-20 °C  |  |
| Storage Comment:    | Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to on  |  |
|                     | month. Avoid repeated freeze-thaw cycles.  |  |