

## Datasheet for ABIN7605614

## anti-COX2 antibody



_				
( )	ve.	rv/	101	Λ

Target:

Quantity:	100 μL	
Target:	COX2	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Monoclonal	
Conjugate:	This COX2 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF),	
	Immunoprecipitation (IP), Immunocytochemistry (ICC), Flow Cytometry (FACS)	
Product Details		
Purpose:	Anti-MTCO2 Rabbit Monoclonal Antibody	
Immunogen:	A synthesized peptide derived from human MTCO2	
Clone:	FHI-13	
Isotype:	IgG	
Characteristics:	Anti-MTCO2 Rabbit Monoclonal Antibody (ABIN7605614). Tested in WB, IHC, ICC/IF, IP, Flow	
	Cytometry applications. This antibody reacts with Human.	
Purification:	Affinity-chromatography	
Target Details		
Tanasta	0000	

COX2

## **Target Details**

raiget Details		
Alternative Name:	MT-CO2 (COX2 Products)	
Background:	Synonyms: Cytochrome c oxidase subunit 2,Cytochrome c oxidase polypeptide II,MT-CO2,COI	
	COXII, MTCO2,	
	Tissue Specificity: Widely expressed with highest levels of expression in liver and heart.	
	Expressed at higher levels in cancer cell lines (e.g. A-549 and HeLa) than in normal cell lines	
	(e.g. HEK293)	
Molecular Weight:	70 kDa	
UniProt:	P00403	
Pathways:	Brown Fat Cell Differentiation, Positive Regulation of fat Cell Differentiation	
Application Details		
Application Notes:	WB 1:500-1:2000	
	IHC 1:50-1:200	
	ICC/IF 1:50-1:200	
	IP 1:20	
	FC 1:50	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Reconstitution:	Restore with deionized water (or equivalent) for reconstitution volume of 1.0 mL	
Concentration:	Lot specific	
Buffer:	Rabbit IgG in phosphate buffered saline, pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 %	
	glycerol, 0.4-0.5 mg/mL BSA.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	
	should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one	
	month. Avoid repeated freeze-thaw cycles.	