

[Go to Product page](#)

Datasheet for ABIN760708  
**anti-CDKN2D antibody (AA 65-166) (Biotin)**

## Overview

Quantity:	100 µL
Target:	CDKN2D
Binding Specificity:	AA 65-166
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CDKN2D antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CDKN2D/p19 INK4d
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat
Purification:	Purified by Protein A.

## Target Details

Target:	CDKN2D
Alternative Name:	CDKN2D ( <a href="#">CDKN2D Products</a> )
Background:	Synonyms: CDK inhibitor p19INK4d, CDKN2D, cell cycle inhibitor, Nur77 associating protein,

## Target Details

Cyclin dependent kinase inhibitor 2D, Cyclin-dependent kinase 4 inhibitor D, Cyclin-dependent kinase inhibitor 2D p19, inhibits CDK4, inhibitor of cyclin-dependent kinase 4d, INK4D, p19, p19-INK4D, p19INK4d, Similar to cyclin-dependent kinase inhibitor 2D, CDN2D\_HUMAN.

Background: The protein encoded by this gene is a member of the INK4 family of cyclin-dependent kinase inhibitors. This protein has been shown to form a stable complex with CDK4 or CDK6, and prevent the activation of the CDK kinases, thus function as a cell growth regulator that controls cell cycle G1 progression. The abundance of the transcript of this gene was found to oscillate in a cell-cycle dependent manner with the lowest expression at mid G1 and a maximal expression during S phase. The negative regulation of the cell cycle involved in this protein was shown to participate in repressing neuronal proliferation, as well as spermatogenesis. Two alternatively spliced variants of this gene, which encode an identical protein, have been reported. [provided by RefSeq, Jul 2008].

Gene ID:	1032
Pathways:	<a href="#">Cell Division Cycle</a> , <a href="#">Sensory Perception of Sound</a> , <a href="#">Mitotic G1-G1/S Phases</a> , <a href="#">Negative Regulation of intrinsic apoptotic Signaling</a>

## Application Details

Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.

## Handling

---

Expiry Date: 12 months