antibodies .- online.com





anti-FCGR2B antibody (pTyr292) (Biotin)



	do to i roddot page

Overview	
Quantity:	100 μL
Target:	FCGR2B
Binding Specificity:	pTyr292
Reactivity:	Human, Mouse, Monkey
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FCGR2B antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)),
	Immunohistochemistry (Frozen Sections) (IHC (fro))
Product Details	
Immunogen:	KLH conjugated synthetic phosphopeptide derived from human CD32B around the
	phosphorylation site of Tyr292.
Isotype:	IgG
Cross-Reactivity:	Human, Monkey, Mouse
Predicted Reactivity:	Rat,Dog,Sheep,Pig,Rabbit,Guinea Pig
Purification:	Purified by Protein A.
Target Details	
Target:	FCGR2B

Target Details

Alternative Name:	CD32B (FCGR2B Products)
Background:	Synonyms: CD32, FCG2, CD32B, FCGR2, IGFR2, Low affinity immunoglobulin gamma Fc region receptor II-b, IgG Fc receptor II-b, CDw32, Fc-gamma RII-b, Fc-gamma-RIIb, FcRII-b, FCGR2B
	Background: Receptor for the Fc region of complexed or aggregated immunoglobulins gamma.
	Low affinity receptor. Involved in a variety of effector and regulatory functions such as
	phagocytosis of immune complexes and modulation of antibody production by B-cells. Binding
	to this receptor results in down-modulation of previous state of cell activation triggered via
	antigen receptors on B-cells (BCR), T-cells (TCR) or via another Fc receptor. Isoform IIB1 fails to
	mediate endocytosis or phagocytosis. Isoform IIB2 does not trigger phagocytosis.
Gene ID:	2213
UniProt:	P31994
Pathways:	Cellular Response to Molecule of Bacterial Origin, Regulation of Leukocyte Mediated Immunity,
	Production of Molecular Mediator of Immune Response, BCR Signaling
Application Details	
Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months