



[Go to Product page](#)

Datasheet for ABIN761505
anti-CPEB1 antibody (AA 451-550) (Cy5)

Overview

| | |
|----------------------|--|
| Quantity: | 100 µL |
| Target: | CPEB1 |
| Binding Specificity: | AA 451-550 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This CPEB1 antibody is conjugated to Cy5 |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| | |
|-----------------------|---|
| Immunogen: | KLH conjugated synthetic peptide derived from human CPEB1 |
| Isotype: | IgG |
| Predicted Reactivity: | Human, Mouse, Rat, Cow, Pig, Horse, Chicken, Rabbit |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|---|
| Target: | CPEB1 |
| Alternative Name: | CPEB1 (CPEB1 Products) |
| Background: | Synonyms: CPE-binding protein 1, CPE-BP1, CPEB 1, CPEB, CPEB1, CPEB1 protein, |

Target Details

CPEB1_HUMAN, Cytoplasmic polyadenylation binding protein 1, Cytoplasmic polyadenylation element binding protein, Cytoplasmic polyadenylation element binding protein 1, Cytoplasmic polyadenylation element-binding protein 1, h-CEBP, hCPEB-1

Background: Sequence-specific RNA-binding protein that regulates mRNA cytoplasmic polyadenylation and translation initiation during oocyte maturation, early development and at postsynapse sites of neurons. Binds to the cytoplasmic polyadenylation element (CPE), an uridine-rich sequence element (consensus sequence 5'-UUUUUUAU-3') within the mRNA 3'-UTR. In absence of phosphorylation and in association with TACC3 is also involved as a repressor of translation of CPE-containing mRNA, a repression that is relieved by phosphorylation or degradation (By similarity). Involved in the transport of CPE-containing mRNA to dendrites, those mRNAs may be transported to dendrites in a translationally dormant form and translationally activated at synapses (By similarity). Its interaction with APLP1 promotes local CPE-containing mRNA polyadenylation and translation activation (By similarity). Induces the assembly of stress granules in the absence of stress.

Gene ID: 64506

Pathways: [Synaptic Membrane](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Handling

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months