# antibodies -online.com





# anti-FGFRL1 antibody (AA 231-330)



Go to Product page

$\sim$					
()	VE	۲۱	/1	$\triangle$	Λ

Quantity:	100 μL	
Target:	FGFRL1	
Binding Specificity:	AA 231-330	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This FGFRL1 antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human FGFR5/FGFRL1	
Isotype:	IgG	
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Horse,Chicken,Rabbit	
Purification:	Purified by Protein A.	

### **Target Details**

Target:	FGFRL1
Alternative Name:	Fgfr5 (FGFRL1 Products)

#### **Target Details**

#### Background:

Synonyms: FGF homologous factor receptor, FGF receptor like protein 1, FGFR 5, FGFR like protein, FGFR5, FGFRL1, FHFR, Fibroblast growth factor receptor 5, Fibroblast growth factor receptor like 1, FGRL1\_HUMAN.

Background: FGFRL1 is a member of the fibroblast growth factor receptor (FGFR) family, where amino acid sequence is highly conserved between members and throughout evolution. FGFR family members differ from one another in their ligand affinities and tissue distribution. A full-length representative protein would consist of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of the protein interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. A marked difference between FGFRL1 and the other family members is its lack of a cytoplasmic tyrosine kinase domain. The result is a transmembrane receptor that could interact with other family members and potentially inhibit signaling. Multiple alternatively spliced transcript variants encoding the same isoform have been found.

Gene ID:

53834

Pathways:

**RTK Signaling** 

#### **Application Details**

Application Notes:

WB 1:300-5000

ELISA 1:500-1000

IHC-P 1:200-400

IHC-F 1:100-500

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin

## Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.	
Expiry Date:	12 months	