antibodies

Datasheet for ABIN762029 anti-MFSD2A antibody (AA 331-430) (Cy3)



Overview

Quantity:	100 µL	
Target:	MFSD2A	
Binding Specificity:	AA 331-430	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This MFSD2A antibody is conjugated to Cy3	
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human MFSD2A	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Predicted Reactivity:	Dog,Cow,Pig,Horse,Chicken,Rabbit	
Purification:	Purified by Protein A.	
Target Details		

Target:	MFSD2A
Alternative Name:	MFSD2A (MFSD2A Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN762029 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

Background:	Synonyms: NLS1, MFSD2, Sodium-dependent lysophosphatidylcholine symporter 1, Sodium-
	dependent LPC symporter 1, Major facilitator superfamily domain-containing protein 2A,
	MFSD2A, HMFN0656, PP9177, UNQ300/PRO341
	Background: Sodium-dependent lysophosphatidylcholine (LPC) symporter, which plays an
	essential role for blood-brain barrier formation and function. Specifically expressed in
	endothelium of the blood-brain barrier of micro-vessels and transports LPC into the brain.
	Transport of LPC is essential because it constitutes the major mechanism by which
	docosahexaenoic acid (DHA), an omega-3 fatty acid that is essential for normal brain growth
	and cognitive function, enters the brain. Transports LPC carrying long-chain fatty acids such
	LPC oleate and LPC palmitate with a minimum acyl chain length of 14 carbons. Does not
	transport docosahexaenoic acid in unesterified fatty acid. Specifically required for blood-brain
	barrier formation and function, probably by mediating lipid transport. Not required for central
	nervous system vascular morphogenesis (By similarity). Acts as a transporter for tunicamycin
	an inhibitor of asparagine-linked glycosylation. In placenta, acts as a receptor for ERVFRD-
	1/syncytin-2 and is required for trophoblast fusion (PubMed:18988732).
Gene ID:	84879
UniProt:	Q8NA29
Application Details	
Application Notes:	FCM 1:20-100
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 an
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN762029 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

lond	lina
land	

	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months