# antibodies .- online.com





anti-HLTF antibody (AA 201-300)



$\sim$			
	N/P	r\/	i⊢₩

Quantity:	100 μL
Target:	HLTF
Binding Specificity:	AA 201-300
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HLTF antibody is un-conjugated
Application:	ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffinembedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human HLTF.
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Pig, Horse, Rabbit
Purification:	Purified by Protein A.
Target Details	

Target:	HLTF
Alternative Name:	HLTF (HLTF Products)

#### **Target Details**

#### Background:

Synonyms: DNA-binding protein/plasminogen activator inhibitor 1 regulator, Helicase like transcription factor, Helicase-like transcription factor, HIP116, HIP116A, HLTF 1, HLTF, HLTF\_HUMAN, HLTF1, p113, RING finger protein 80, RNF80, SMARC A3, SMARCA 3, SMARCA3, SNF2-like 3, SNF2L3, Sucrose nonfermenting protein 2 like 3, Sucrose nonfermenting protein 2-like 3, SWI/SNF related matrix associated actin dependent regulator of chromatin subfamily A member 3, SWI/SNF-related matrix-associated actin-dependent regulator of chromatin a3, SWI/SNF-related matrix-associated actin-dependent regulator of chromatin subfamily A member 3, ZBU 1, ZBU1.

Background: Has both helicasee and E3 ubiquitin ligase activities. Possesses intrinsic ATP-dependent nucleosome-remodeling activity, This activity may be required for transcriptional activation or repression of specific target promoters (By similarity). These may include the SERPINE1 and HIV-1 promoters and the SV40 enhancer, to which this protein can bind directly. Plays a role in error-free postreplication repair (PRR) of damaged DNA and maintains genomic stability through acting as a ubiquitin ligase for 'Lys-63'-linked polyubiquitination of chromatin-bound PCNA.

Gene ID:

6596

## **Application Details**

Application Notes:	

ELISA 1:500-1000

IHC-P 1:200-400

IHC-F 1:100-500

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

# Handling

Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months