antibodies .- online.com





anti-HLTF antibody (AA 201-300) (Cy3)



Overview

Quantity:	100 μL	
Target:	HLTF	
Binding Specificity:	AA 201-300	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This HLTF antibody is conjugated to Cy3	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human HLTF.
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	HLTF
Alternative Name:	HLTF (HLTF Products)
Background:	Synonyms: DNA-binding protein/plasminogen activator inhibitor 1 regulator, Helicase like

transcription factor, Helicase-like transcription factor, HIP116, HIP116A, HLTF 1, HLTF, HLTF_HUMAN, HLTF1, p113, RING finger protein 80, RNF80, SMARC A3, SMARCA 3, SMARCA3, SNF2-like 3, SNF2L3, Sucrose nonfermenting protein 2 like 3, Sucrose nonfermenting protein 2-like 3, SWI/SNF related matrix associated actin dependent regulator of chromatin subfamily A member 3, SWI/SNF-related matrix-associated actin-dependent regulator of chromatin a3, SWI/SNF-related matrix-associated actin-dependent regulator of chromatin subfamily A member 3, ZBU 1, ZBU1.

Background: Has both helicasee and E3 ubiquitin ligase activities. Possesses intrinsic ATP-dependent nucleosome-remodeling activity, This activity may be required for transcriptional activation or repression of specific target promoters (By similarity). These may include the SERPINE1 and HIV-1 promoters and the SV40 enhancer, to which this protein can bind directly. Plays a role in error-free postreplication repair (PRR) of damaged DNA and maintains genomic stability through acting as a ubiquitin ligase for 'Lys-63'-linked polyubiquitination of chromatin-bound PCNA.

Gene ID:

6596

Application Details

IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

\vdash	land	linc
	iaria	11116

Expiry Date:

12 months