

Datasheet for ABIN7625776 anti-PTPMT1 antibody (AA 28-201)



()	V		rV	ĺ	9	V	V
'	\mathcal{I}	٧V	<u> </u>	v	1	$\overline{}$	٧	٧

Quantity:	100 μg		
Target:	PTPMT1		
Binding Specificity:	AA 28-201		
Reactivity:	Human		
Host:	Rabbit		
Clonality:	Polyclonal		
Conjugate:	This PTPMT1 antibody is un-conjugated		
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (IHC)		
Product Details			
Dumana	DTDMT1 Debut land Autiliards		
Purpose:	PTPMT1 Polyclonal Antibody		
Immunogen:	Recombinant Human Phosphatidylglycerophosphatase protein-tyrosine phosphatase 1 protein		
•			
Immunogen:	Recombinant Human Phosphatidylglycerophosphatase protein-tyrosine phosphatase 1 protein		
Immunogen: Isotype:	Recombinant Human Phosphatidylglycerophosphatase protein-tyrosine phosphatase 1 protein IgG		
Immunogen: Isotype:	Recombinant Human Phosphatidylglycerophosphatase protein-tyrosine phosphatase 1 protein IgG The PTPMT1 Polyclonal Antibody (Species: Human) has been validated for the following		
Immunogen: Isotype: Characteristics:	Recombinant Human Phosphatidylglycerophosphatase protein-tyrosine phosphatase 1 protein IgG The PTPMT1 Polyclonal Antibody (Species: Human) has been validated for the following applications: ELISA, WB, IHC.		
Immunogen: Isotype: Characteristics: Purification:	Recombinant Human Phosphatidylglycerophosphatase protein-tyrosine phosphatase 1 protein IgG The PTPMT1 Polyclonal Antibody (Species: Human) has been validated for the following applications: ELISA, WB, IHC. Protein G purified		

Target Details

Storage:

Expiry Date:

Storage Comment:

Alternative Name:	PTPMT1 (PTPMT1 Products)		
UniProt:	Q8WUK0		
Pathways:	Inositol Metabolic Process		

Application Details	
Application Notes: Optimal working dilution should be determined by the investigator.	
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	Lot specific
Buffer:	0.03 % Proclin 300, 50 % glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

 2°C to 8°C for frequent use, -20°C for 12 months. Avoid repeated freeze/thaw cycles.

4 °C,-20 °C

12 months