antibodies - online.com





Datasheet for ABIN762581

anti-SPARCL1 antibody (AA 601-664)



Publication



()	1 /	-	rv	1 /	71	A
	1//	\vdash	1 \/	16		۱/۱
\sim	٧.	\sim	1 V	ı١	_	V١

Quantity:	100 μL
Target:	SPARCL1
Binding Specificity:	AA 601-664
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SPARCL1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human SPARCL1
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

|--|

Target Details

Alternative Name:	Sparcl1 (SPARCL1 Products)	
Background:	Synonyms: Ecm2, RAGS1 antigen, Extracellular matrix protein 2, Hevin, High endothelial venule	
	protein, MAST 9, mast9, Matrix glycoprotein Sc1, PIG33, proliferation inducing protein 33, SC1,	
	SPARC like 1, SPARC-like protein 1, Sparcl1, SPRL1_HUMAN.	
	Background: SPARC (secreted protein acidic and rich in cysteine) is a phosphorylated, acidic,	
	glycine-rich glycoprotein that is secreted by endothelial cells and is present in large amounts in	
	the parietal endoderm of mouse embryos and in human placenta. SPARC-like protein 1	
	(SPARCL1), also known as high endothelial venule protein (Hevin) or MAST9, is a 664 amino	
	acid member of the SPARC family of proteins. Highly expressed in lymph node, heart, lung,	
	brain, skeletal muscle, ovary, colon and small intestine, SPARCL1 is a secreted protein that	
	contains one EF-hand domain, one follistatin-like domain and one Kazal-like domain. SPARCL1	
	is implicated to play a role in neuronal remodeling and tumor suppression. The gene encoding	
	SPARCL1 maps to chromosome 4q22.1.	
Gene ID:	8404	
Application Details		
Application Notes:	WB 1:300-5000	
	ELISA 1:500-1000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	

Handling

Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months
Publications	
Product cited in:	Lai, Wu, Zou, Xie, Zhang, Zhao, Zhao, Wang, Ji: "Secretome Analyses of Aβ(1-42) Stimulated
	Hippocampal Astrocytes Reveal that CXCL10 is Involved in Astrocyte Migration." in: Journal of
	proteome research, (2013) (PubMed).