

Datasheet for ABIN762851  
**anti-WNT8A antibody (AA 201-300)**[Go to Product page](#)

## 1 Image

## Overview

Quantity:	100 µL
Target:	WNT8A
Binding Specificity:	AA 201-300
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This WNT8A antibody is un-conjugated
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Wnt8A
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Sheep,Horse
Purification:	Purified by Protein A.

## Target Details

Target:	WNT8A
---------	-------

## Target Details


Alternative Name:	Wnt8a ( <a href="#">WNT8A Products</a> )
Background:	Synonyms: WNT8D, Protein Wnt-8a, Protein Wnt-8d, WNT8A Background: Ligand for members of the frizzled family of seven transmembrane receptors. May play an important role in the development and differentiation of certain forebrain structures, notably the hippocampus.
Gene ID:	7478
UniProt:	<a href="#">Q9H1J5</a>
Pathways:	<a href="#">WNT Signaling</a>

## Application Details

Application Notes:	WB 1:300-5000 ELISA 1:500-1000 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months



#### Immunohistochemistry

**Image 1.** Formalin-fixed and paraffin embedded rat brain labeled with Anti-Wnt8A Polyclonal Antibody, Unconjugated (ABIN762851) at 1:200 followed by conjugation to the secondary antibody and DAB staining