



[Go to Product page](#)

Datasheet for ABIN762991
anti-KDM5B antibody (AA 65-165) (Cy5.5)

Overview

Quantity:	100 µL
Target:	KDM5B
Binding Specificity:	AA 65-165
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This KDM5B antibody is conjugated to Cy5.5
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human KDM5B/Jarid1B
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Predicted Reactivity:	Mouse,Dog,Cow,Pig,Horse,Chicken,Guinea Pig
Purification:	Purified by Protein A.

Target Details

Target:	KDM5B
Alternative Name:	KDM5B (KDM5B Products)

Target Details

Background: Synonyms: CT31, PLU1, PUT1, PLU-1, JARID1B, PPP1R98, RBBP2H1A, Lysine-specific demethylase 5B, Cancer/testis antigen 31, Histone demethylase JARID1B, Jumonji/ARID domain-containing protein 1B, Retinoblastoma-binding protein 2 homolog 1, RBP2-H1, KDM5B, RBBP2H1

Background: Histone demethylase that demethylates 'Lys-4' of histone H3, thereby playing a central role in histone code. Does not demethylate histone H3 'Lys-9' or H3 'Lys-27'. Demethylates trimethylated, dimethylated and monomethylated H3 'Lys-4'. Acts as a transcriptional corepressor for FOXG1B and PAX9. Favors the proliferation of breast cancer cells by repressing tumor suppressor genes such as BRCA1 and HOXA5. In contrast, may act as a tumor suppressor for melanoma. Represses the CLOCK-ARNTL/BMAL1 heterodimer-mediated transcriptional activation of the core clock component PER2 (By similarity).

Gene ID: 10765

UniProt: [Q9UGL1](#)

Pathways: [Warburg Effect](#)

Application Details

Application Notes: FCM 1:20-100
IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Handling

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months