

Datasheet for ABIN7630066

Recombinant anti-Coronavirus Spike Glycoprotein antibody



Overview	
Quantity:	100 μL
Target:	Coronavirus Spike Glycoprotein (CoV S)
Reactivity:	Various Species
Host:	Mouse
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This Coronavirus Spike Glycoprotein antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunoprecipitation (IP), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Flow Cytometry (FACS)
Product Details	
Purpose:	Recombinant Antibody to Spike Protein (SP)
Isotype:	IgG
Specificity:	The antibody is a mouse monoclonal antibody raised against SP. It has been selected for its ability to recognize SP in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	
Target:	Coronavirus Spike Glycoprotein (CoV S)
Alternative Name:	Spike Protein (CoV S Products)

Target Details

Target Type:	Viral Protein
Background:	S Protein
UniProt:	P0DTC2

Application Details

Application Notes:	Western blotting: 0.2-2 μg/mL,1:500-5000 Immunohistochemistry: 5-20 μg/mL,1:50-200
	Immunocytochemistry: 5-20 μg/mL,1:50-200 Optimal working dilutions must be determined by
	end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.