

Datasheet for ABIN7630915 **anti-CXCL3 antibody (Biotin)**



[Go to Product page](#)

Overview

Quantity:	1 mL
Target:	CXCL3
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CXCL3 antibody is conjugated to Biotin
Application:	Immunohistochemistry (IHC), Western Blotting (WB), Immunocytochemistry (ICC)

Product Details

Purpose:	Biotin-Linked Polyclonal Antibody to Chemokine (C-X-C Motif) Ligand 3 (CXCL3)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against CXCL3. It has been selected for its ability to recognize CXCL3 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	CXCL3
Alternative Name:	Chemokine (C-X-C Motif) Ligand 3 (CXCL3 Products)
Background:	GROg, SCYB3, GRO3, GRO-G, MIP2-B, MIP2b, CINC2b, CINC2b, Growth Regulated Oncogene Gamma, Macrophage inflammatory protein 2-beta

Target Details

UniProt: [P19876](#)

Pathways: [Cellular Response to Molecule of Bacterial Origin, Autophagy](#)

Application Details

Application Notes: Western blotting: 0.2-2 µg/mL, 1:250-2500 Immunohistochemistry: 5-20 µg/mL, 1:25-100
Immunocytochemistry: 5-20 µg/mL, 1:25-100 Optimal working dilutions must be determined by end user.

Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 500 µg/mL

Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.