

Datasheet for ABIN7631093

anti-Corticosterone antibody (Biotin)



Overview

Quantity:	1 mL
Target:	Corticosterone (CORT)
Reactivity:	Various Species
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Corticosterone antibody is conjugated to Biotin
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Product Details	
Purpose:	Biotin-Linked Polyclonal Antibody to Corticosterone (Cort)
Immunogen:	PAA540Ge01Polyclonal Antibody to Corticosterone (Cort)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against Cort. It has been selected for its ability to recognize Cort in immunohistochemical staining and western blotting.
Cross-Reactivity:	Various Species
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	
Target:	Corticosterone (CORT)
Alternative Name:	Corticosterone (CORT Products)

Target Details

Target Type:	Hormone
Background:	Corticosrone
Application Details	
Application Notes:	Immunocytochemistry in formalin fixed cells: 5-20 μ g/mL Immunohistochemistry in formalin fixed frozen section: 5-20 μ g/mL Immunohistochemistry in paraffin section: 5-20 μ g/mL Enzyme-linked Immunosorbent Assay: 0.05-2 μ g/mL Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	500 μg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.