

Datasheet for ABIN7631788 **anti-HSPG antibody (FITC)**



[Go to Product page](#)

Overview

Quantity:	1 mL
Target:	HSPG
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HSPG antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Purpose:	FITC-Linked Polyclonal Antibody to Heparan Sulfate Proteoglycan (HSPG)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against HSPG. It has been selected for its ability to recognize HSPG in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	HSPG
Alternative Name:	Heparan Sulfate Proteoglycan (HSPG Products)
Background:	SDC2, HSPG1, SYND2, Fibroglycan, Syndecan 2, Syndecan Proteoglycan 2, Heparan Sulfate Proteoglycan 1, Cell Surface-Associated, Heparan Sulphate Proteoglycan

Target Details

UniProt: [P34741](#)

Application Details

Application Notes: Western blotting: 0.2-2 µg/mL, 1:250-2500 Immunohistochemistry: 5-20 µg/mL, 1:25-100
Immunocytochemistry: 5-20 µg/mL, 1:25-100 Optimal working dilutions must be determined by end user.

Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 500 µg/mL

Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.