

Datasheet for ABIN763213

anti-RAC1 antibody (AA 101-192) (Biotin)



Go to Product page

_					
	1//	r	Vİ	\triangle	۸/
	V		VI		/ V

Quantity:	100 μL	
Target:	RAC1	
Binding Specificity:	AA 101-192	
Reactivity:	Human, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This RAC1 antibody is conjugated to Biotin	
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry	
	(Frozen Sections) (IHC (fro))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human RAC1/RAC2.	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse	
Cross-Reactivity: Predicted Reactivity:	Human, Mouse Rat,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit,Guinea Pig	
Predicted Reactivity:	Rat,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit,Guinea Pig	
Predicted Reactivity: Purification:	Rat,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit,Guinea Pig	

Target Details

<u> </u>			
Background:	Synonyms: MIG5, p21 Rac1, Rac 1, Rac-1, p21 Rac2, PAK1, Ras like protein TC25, Ras related		
	C3 botulinum toxin substrate 1, Ras related C3 botulinum toxin substrate 2, RAC1_HUMAN,		
	RAC2_HUMAN.		
	Background: The protein encoded by this gene is a GTPase which belongs to the RAS		
	superfamily of small GTP-binding proteins. Members of this superfamily appear to regulate a		
	diverse array of cellular events, including the control of cell growth, cytoskeletal reorganization,		
	and the activation of protein kinases. Two transcript variants encoding different isoforms have		
	been found for this gene. [provided by RefSeq, Mar 2009].		
Gene ID:	5879, 5880		
Pathways:	WNT Signaling, Regulation of Actin Filament Polymerization, Cell-Cell Junction Organization,		
	Thromboxane A2 Receptor Signaling, VEGF Signaling		
Application Details			
Application Notes:	IHC-P 1:200-400		
	IHC-F 1:100-500		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
	50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
	handled by trained staff only.		
Storage:	-20 °C		
Storage Comment:	Store at -20°C for 12 months.		
Expiry Date:	12 months		
Expiry Date:	12 months		