

## Datasheet for ABIN7632157

## anti-IL5RA antibody (Biotin)



## Overview

| Overview          |  |
|-------------------|--|
| Quantity:         | 1 mL   |
| Target:           | IL5RA  |
| Reactivity:       | Human  |
| Host:             | Rabbit   |
| Clonality:        | Polyclonal   |
| Conjugate:        | This IL5RA antibody is conjugated to Biotin  |
| Application:      | Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)   |
| Product Details   |  |
| Purpose:          | Biotin-Linked Polyclonal Antibody to Interleukin 5 Receptor Alpha (IL5Ra)  |
| Immunogen:        | PAB510Hu01Polyclonal Antibody to Interleukin 5 Receptor Alpha (IL5Ra)  |
| Isotype:          | IgG  |
| Specificity:      | The antibody is a rabbit polyclonal antibody raised against IL5Ra. It has been selected for its ability to recognize IL5Ra in immunohistochemical staining and western blotting. |
| Purification:     | Antigen-specific affinity chromatography followed by Protein A affinity chromatography   |
| Target Details    |  |
| Target:           | IL5RA  |
| Alternative Name: | Interleukin 5 Receptor Alpha (IL5RA Products)  |
| Background:       | CDw125, CD125, IL5R-A, IL5-RA, HSIL5R3, IL5R, IL5R-A   |
|                   |  |

## **Target Details**

| UniProt:              | Q01344   |
|-----------------------|--|
| Pathways:             | JAK-STAT Signaling   |
| Application Details   |  |
| - Application Details |  |
| Application Notes:    | Western blotting: 0.5-2 $\mu$ g/mL Immunocytochemistry in formalin fixed cells: 5-20 $\mu$ g/mL  |
|                       | Immunohistochemistry in formalin fixed frozen section: 5-20 µg/mL Immunohistochemistry in        |
|                       | paraffin section: 5-20 μg/mL Optimal working dilutions must be determined by end user.           |
| Comment:              | The thermal stability is described by the loss rate. The loss rate was determined by accelerated |
|                       | thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious          |
|                       | degradation and precipitation were observed. The loss rate is less than 5% within the expiration |
|                       | date under appropriate storage condition.  |
|                       |  |
| Restrictions:         | For Research Use only  |
|                       |  |
| Handling              |  |
| Format:               | Liquid   |
| Concentration:        | 500 μg/mL  |
| Buffer:               | 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.                                 |
| Preservative:         | ProClin  |
| Precaution of Use:    | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be               |
|                       | handled by trained staff only.   |
| 04                    | 4.00.00.00   |
| Storage:              | 4 °C,-20 °C  |
| Storage Comment:      | Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without  |
|                       | detectable loss of activity. Avoid repeated freeze-thaw cycles.                                  |
|                       |  |