

Datasheet for ABIN7632213

anti-IDH1 antibody (Biotin)



Overview

Overview	
Quantity:	1 mL
Target:	IDH1
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This IDH1 antibody is conjugated to Biotin
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Product Details	
Purpose:	Biotin-Linked Monoclonal Antibody to Isocitrate Dehydrogenase 1, Soluble (IDH1)
Immunogen:	MAH839Hu26Monoclonal Antibody to Isocitrate Dehydrogenase 1, Soluble (IDH1)
Clone:	C3
Isotype:	IgG
Specificity:	The antibody is a mouse monoclonal antibody raised against IDH1. It has been selected for its
	ability to recognize IDH1 in immunohistochemical staining and western blotting.
Purification:	Protein A + Protein G affinity chromatography
Target Details	
Target:	IDH1
Alternative Name:	Isocitrate Dehydrogenase 1, Soluble (IDH1 Products)

Target Details

Target Details	
Background:	PICD, IDP, Oxalosuccinate decarboxylase, NADP(+)-specific ICDH, Cytosolic NADP-isocitrate dehydrogenase
	denydrogenase
UniProt:	075874
Pathways:	Warburg Effect
Application Details	
Application Notes:	Western blotting: $0.5-2~\mu g/mL$ Immunocytochemistry in formalin fixed cells: $5-20~\mu g/mL$ Immunohistochemistry in formalin fixed frozen section: $5-20~\mu g/mL$ Immunohistochemistry in paraffin section: $5-20~\mu g/mL$ Enzyme-linked Immunosorbent Assay: $0.05-2~\mu g/mL$ Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.