

## Datasheet for ABIN7632298

## anti-Kininogen (HMW) antibody (FITC)



()	ve	r\/i	Δ	۱۸/
$\circ$	V C	1 V		v v

Quantity:	1 mL
Target:	Kininogen (HMW)
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Kininogen (HMW) antibody is conjugated to FITC
Application:	Immunohistochemistry (IHC), Western Blotting (WB), Immunocytochemistry (ICC), Immunofluorescence (IF)

## **Product Details**

Alternative Name:

Background:

Purpose:	FITC-Linked Polyclonal Antibody to High Molecular Weight Kininogen (HMWK)	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against HMWK. It has been selected for its ability to recognize HMWK in immunohistochemical staining and western blotting.	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography	
Target Details		
Target:	Kininogen (HMW)	

High Molecular Weight Kininogen (Kininogen (HMW) Products)

Williams-Fitzgerald-Flaujeac Factor, Fitzgerald Factor, HMWK-Kallikrein Factor

## **Target Details**

UniProt:	P08934	
Application Details		
Application Notes:	Western blotting: 0.2-2 μg/mL,1:250-2500 Immunohistochemistry: 5-20 μg/mL,1:25-100 Immunocytochemistry: 5-20 μg/mL,1:25-100 Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	500 μg/mL	
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.	