

## Datasheet for ABIN7632617 anti-MYH8 antibody (FITC)



## Overview

Quantity:	1 mL
Target:	MYH8
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MYH8 antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunofluorescence (IF)

## **Product Details**

FIOUUCI Details	
Purpose:	FITC-Linked Polyclonal Antibody to Myosin Heavy Chain 8, Skeletal Muscle, Perinatal (MYH8)
Immunogen:	PAD421Ra01Polyclonal Antibody to Myosin Heavy Chain 8, Skeletal Muscle, Perinatal (MYH8)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against MYH8. It has been selected for its ability to recognize MYH8 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	

Target:	MYH8
Alternative Name:	Myosin Heavy Chain 8, Skeletal Muscle, Perinatal (MYH8 Products)

## **Target Details**

Background:	MyHC-peri, MyHC-pn, MyHC-perinatal, Myosin heavy chain, skeletal muscle, perinatal
UniProt:	P04462
Application Details	
Application Notes:	Western blotting: 0.5-2 μg/mL Immunohistochemistry: 5-20 μg/mL Immunocytochemistry: 5-20 μg/mL Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	500 μg/mL
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.