

Datasheet for ABIN7632978
anti-POMC antibody (FITC)



[Go to Product page](#)

Overview

Quantity:	1 mL
Target:	POMC
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This POMC antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Purpose:	FITC-Linked Polyclonal Antibody to Proopiomelanocortin (POMC)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against POMC. It has been selected for its ability to recognize POMC in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	POMC
Alternative Name:	Proopiomelanocortin (POMC Products)
Background:	POC, ProOMC, Pro-OMC, Pro Opiomelanocortin, Corticotropin-lipotropin

Target Details

UniProt:	P01189
Pathways:	Metabolism of Steroid Hormones and Vitamin D , Peptide Hormone Metabolism , Hormone Activity , Feeding Behaviour

Application Details

Application Notes:	Western blotting: 0.2-2 µg/mL, 1:250-2500 Immunohistochemistry: 5-20 µg/mL, 1:25-100 Immunocytochemistry: 5-20 µg/mL, 1:25-100 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	500 µg/mL
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.