

Datasheet for ABIN7633017 **anti-PTPRS antibody (FITC)**



[Go to Product page](#)

Overview

Quantity:	1 mL
Target:	PTPRS
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PTPRS antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunofluorescence (IF)

Product Details

Purpose:	FITC-Linked Polyclonal Antibody to Protein Tyrosine Phosphatase Receptor Type S (PTPRS)
Immunogen:	PAB295Mu01 Polyclonal Antibody to Protein Tyrosine Phosphatase Receptor Type S (PTPRS)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against PTPRS. It has been selected for its ability to recognize PTPRS in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	PTPRS
Alternative Name:	Protein Tyrosine Phosphatase Receptor Type S (PTPRS Products)

Target Details

Background:	PTPR-S, PTPSIGMA, PTP SIGMA, Receptor-type tyrosine-protein phosphatase sigma
UniProt:	B0V2N1

Application Details

Application Notes:	Western blotting: 0.5-2 µg/mL Immunohistochemistry: 5-20 µg/mL Immunocytochemistry: 5-20 µg/mL Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	500 µg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.