

Datasheet for ABIN7633038

anti-REG3g antibody (Biotin)



()	ver	~\ /I	~ 1 A
	$\vee \square$	\/ I	ᆸᇄ
\sim	v Cı	V I	$\smile v$

Quantity:	1 mL
Target:	REG3g
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This REG3g antibody is conjugated to Biotin
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Product Details	
Purpose:	Biotin-Linked Polyclonal Antibody to Regenerating Islet Derived Protein 3 Gamma (REG3g)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against REG3g. It has been selected for its ability to recognize REG3g in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	
Target:	REG3g
Alternative Name:	Regenerating Islet Derived Protein 3 Gamma (REG3g Products)
Background:	REG-III, PAP1B, PAPIB, UNQ429, Pancreatitis-associated protein 1B, Regenerating islet-derived protein III-gamma

Target Details

UniProt:	009049	
Pathways:	Activation of Innate immune Response	
Application Details		
Application Notes:	Western blotting: 0.2-2 μ g/mL,1:250-2500 Immunohistochemistry: 5-20 μ g/mL,1:25-100 Immunocytochemistry: 5-20 μ g/mL,1:25-100 Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	500 μg/mL	
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.	