antibodies -online.com





Datasheet for ABIN763335

anti-ARL3 antibody (AA 1-100) (Cy5)

Go to Product page

| <i>ا</i> ۱ | 1 | - | M | /1 | 0 | |
|------------|-----|----------|-----|-----|--------|------|
| u | 1// | \vdash | 1 \ | / I | - | \/\/ |
| \sim | ٧. | \sim | r٧ | | \sim | v v |

| Quantity: | 100 μL |
|----------------------|---------------------------------------------------------------------------------------------------------|
| Target: | ARL3 |
| Binding Specificity: | AA 1-100 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This ARL3 antibody is conjugated to Cy5 |
| Application: | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human ARL3 |
|-----------------------|----------------------------------------------------------|
| Isotype: | IgG |
| Predicted Reactivity: | Human,Mouse,Rat,Cow,Pig,Horse,Rabbit |
| Purification: | Purified by Protein A. |

Target Details

| Target: | ARL3 |
|---------------------------------------------------------------------------------------------------|----------------------|
| Alternative Name: | ARL3 (ARL3 Products) |
| Background: Synonyms: ADP ribosylation factor like 3, ADP-ribosylation factor-like protein 3, Arf | |

Expiry Date:

12 months

| Target Details | |
|---------------------|--------------------------------------------------------------------------------------------------------------------|
| | protein 3, ARFL3, Arl3, ARL3_HUMAN. |
| | Background: Small GTP-binding protein which cycles between an inactive GDP-bound and an |
| | active GTP-bound form, and the rate of cycling is regulated by guanine nucleotide exchange |
| | factors (GEF) and GTPase-activating proteins (GAP). GTP-binding protein that does not act as |
| | an allosteric activator of the cholera toxin catalytic subunit. Requires assistance from GTPase- |
| | activating proteins (GAPs) like RP2 and PDE6D, in order to cycle between inactive GDP-bound |
| | and active GTP-bound forms. Required for normal cytokinesis and cilia signaling. |
| Gene ID: | 403 |
| Application Details | |
| Application Notes: | IF(IHC-P) 1:50-200 |
| | IF(IHC-F) 1:50-200 |
| | IF(ICC) 1:50-200 |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and |
| | 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |