

Datasheet for ABIN7633547  
**anti-Transferrin antibody (FITC)**



[Go to Product page](#)

## Overview

|              |                                                                                                          |
|--------------|----------------------------------------------------------------------------------------------------------|
| Quantity:    | 1 mL                                                                                                     |
| Target:      | Transferrin (TF)                                                                                         |
| Reactivity:  | Mouse                                                                                                    |
| Host:        | Rabbit                                                                                                   |
| Clonality:   | Polyclonal                                                                                               |
| Conjugate:   | This Transferrin antibody is conjugated to FITC                                                          |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC),<br>Immunofluorescence (IF) |

## Product Details

|               |                                                                                                                                                                            |
|---------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Purpose:      | FITC-Linked Polyclonal Antibody to Transferrin (TF)                                                                                                                        |
| Isotype:      | IgG                                                                                                                                                                        |
| Specificity:  | The antibody is a rabbit polyclonal antibody raised against TF. It has been selected for its ability to recognize TF in immunohistochemical staining and western blotting. |
| Purification: | Antigen-specific affinity chromatography followed by Protein A affinity chromatography                                                                                     |

## Target Details

|                   |                                                                   |
|-------------------|-------------------------------------------------------------------|
| Target:           | Transferrin (TF)                                                  |
| Alternative Name: | Transferrin ( <a href="#">TF Products</a> )                       |
| Background:       | TRF, Siderophilin, Serotransferrin, Beta-1 metal-binding globulin |

## Target Details

|           |                                                  |
|-----------|--------------------------------------------------|
| UniProt:  | <a href="#">Q92111</a>                           |
| Pathways: | <a href="#">Transition Metal Ion Homeostasis</a> |

## Application Details

|                    |                                                                                                                                                                                                                                                                                                                                     |
|--------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Application Notes: | Western blotting: 0.2-2 µg/mL, 1:250-2500 Immunohistochemistry: 5-20 µg/mL, 1:25-100<br>Immunocytochemistry: 5-20 µg/mL, 1:25-100 Optimal working dilutions must be determined by end user.                                                                                                                                         |
| Comment:           | The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition. |
| Restrictions:      | For Research Use only                                                                                                                                                                                                                                                                                                               |

## Handling

|                    |                                                                                                                                                                 |
|--------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Format:            | Liquid                                                                                                                                                          |
| Concentration:     | 500 µg/mL                                                                                                                                                       |
| Buffer:            | PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.                                                                                                     |
| Preservative:      | Sodium azide                                                                                                                                                    |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.                                          |
| Storage:           | 4 °C, -20 °C                                                                                                                                                    |
| Storage Comment:   | Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles. |