

Datasheet for ABIN7633750 anti-Vitamin B12 antibody (FITC)



Overview

Quantity:	1 mL
Target:	Vitamin B12
Reactivity:	Various Species
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Vitamin B12 antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunofluorescence (IF)

Product Details

Target:

Froduct Details	
Purpose:	FITC-Linked Polyclonal Antibody to Cyanocobalamin (CNCbl)
Immunogen:	PAA924Ge01Polyclonal Antibody to Cyanocobalamin (CNCbl)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against CNCbl. It has been selected for its ability to recognize CNCbl in immunohistochemical staining and western blotting.
Cross-Reactivity:	Various Species
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	

Vitamin B12

Target Details

Cyanocobalamin (Vitamin B12 Products)
Chemical
VB12, Vitamin B12, Cobalamin, α-5,6-Dimethylbenzimidazolyl Cobamidcyanide
Immunocytochemistry in formalin fixed cells: 5-20 μ g/mL Immunohistochemistry in formalin fixed frozen section: 5-20 μ g/mL Immunohistochemistry in paraffin section: 5-20 μ g/mL Enzyme-linked Immunosorbent Assay: 0.05-2 μ g/mL Optimal working dilutions must be determined by end user.
The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
For Research Use only
Liquid
500 μg/mL
0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
ProClin
This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
4 °C,-20 °C