

Datasheet for ABIN7633981 anti-Actin, gamma 1 antibody (APC)



Go to Product page

| _ | | | |
|-----|-----|----|----|
| () | V/C | rv | ٨/ |
| | | | |

| Quantity: | 100 μL |
|--------------|--|
| Target: | Actin, gamma 1 (ACTG1) |
| Reactivity: | Human |
| Host: | Guinea Pig |
| Clonality: | Polyclonal |
| Conjugate: | This Actin, gamma 1 antibody is conjugated to APC |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC) |

Product Details

| 1 Toddot Detaile | |
|------------------|---|
| Purpose: | Polyclonal Antibody to Gamma Actin (ACTG) |
| Immunogen: | PAB391Hu52Polyclonal Antibody to Gamma Actin (ACTG) |
| Isotype: | IgG |
| Specificity: | The antibody is a cavia polyclonal antibody raised against ACTG. It has been selected for its ability to recognize ACTG in immunohistochemical staining and western blotting. |
| Purification: | Antigen-specific affinity chromatography followed by Protein A affinity chromatography |
| Target Details | |

| Target: | Actin, gamma 1 (ACTG1) |
|-------------------|------------------------|
| Alternative Name: | ACTG (ACTG1 Products) |

Target Details

| Target Details | | |
|---------------------|--|--|
| Background: | ACT, ACTB, ACTg1, DFNA20, DFNA26, deafness, autosomal dominant 20, deafness, autosomal dominant 26, Actin, cytoplasmic 2 | |
| | dominant 26, Actin, cytopiasmic 2 | |
| UniProt: | P63261 | |
| Pathways: | Myometrial Relaxation and Contraction, Cell-Cell Junction Organization | |
| Application Details | | |
| Application Notes: | Western blotting: 0.5-2 μg/mL, Immunohistochemistry: 5-20 μg/mL, Immunocytochemistry: 5- | |
| | 20 μg/mL,Optimal working dilutions must be determined by end user. | |
| Comment: | The thermal stability is described by the loss rate. The loss rate was determined by accelerated | |
| | thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious | |
| | degradation and precipitation were observed. The loss rate is less than 5% within the expiration | |
| | date under appropriate storage condition. | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 500 μg/mL | |
| Buffer: | 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be | |
| | handled by trained staff only. | |
| Storage: | 4 °C,-20 °C | |
| Storage Comment: | Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without | |
| | detectable loss of activity. Avoid repeated freeze-thaw cycles. | |