

Datasheet for ABIN7634040

anti-ACOX3 antibody



Overview

Quantity:	100 μL
Target:	ACOX3
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ACOX3 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Western Blotting (WB), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Acyl Coenzyme A Oxidase 3, Pristanoyl (ACOX3)
Immunogen:	RPD667Mu01Recombinant Acyl Coenzyme A Oxidase 3, Pristanoyl (ACOX3)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against ACOX3. It has been selected for its ability to recognize ACOX3 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Rat
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	
Target:	ACOX3

Target Details	
Alternative Name:	ACOX3 (ACOX3 Products)
Background:	BRCOX, PRCOX, BRCACox, Branched-chain acyl-CoA oxidase, Pristanoyl-CoA oxidase, Peroxisomal acyl-coenzyme A oxidase 3
UniProt:	Q9EPL9
Pathways:	Monocarboxylic Acid Catabolic Process
Application Details	
Application Notes:	Western blotting: 0.01-2 μg/mL,Immunohistochemistry: 5-20 μg/mL,Immunocytochemistry: 5-20 μg/mL,Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	

Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.