

Datasheet for ABIN7634044
anti-ACSM1 antibody



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	ACSM1
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ACSM1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Purpose:	Polyclonal Antibody to Acyl Coenzyme A Synthetase Medium Chain Family, Member 1 (ACSM1)
Immunogen:	RPD669Hu01Recombinant Acyl Coenzyme A Synthetase Medium Chain Family, Member 1 (ACSM1)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against ACSM1. It has been selected for its ability to recognize ACSM1 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse, Pig, Rat
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	ACSM1
---------	-------

Target Details

Alternative Name:	ACSM1 (ACSM1 Products)
Background:	MACS1, BUCS1, LAE, Butyryl Coenzyme A Synthetase 1, Acyl-CoA synthetase medium-chain family member 1, Middle-chain acyl-CoA synthetase 1, Lipoate-activating enzyme
UniProt:	Q08AH1

Application Details

Application Notes:	Western blotting: 0.01-2 µg/mL, immunohistochemistry: 5-20 µg/mL, Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.