

Datasheet for ABIN7634422 **anti-Fetuin A antibody**

[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	Fetuin A (AHSG)
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Fetuin A antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Alpha-2-Heremans Schmid Glycoprotein (AHSG)
Immunogen:	RPA178Mu02Recombinant Alpha2Heremans Schmid Glycoprotein (AHSG)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against AHSG. It has been selected for its ability to recognize AHSG in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	Fetuin A (AHSG)
Alternative Name:	AHSG (AHSG Products)

Target Details

Background: AHSB, HSBa, A2HS, AHS, FETUA, HSB-A2, Fetuin-A, Alpha 2-HS Glycoprotein, Fetuin A, Alpha-2-Z-Globulin, Ba-alpha-2-Glycoprotein

UniProt: [P29699](#)

Application Details

Application Notes: Western blotting: 0.5-2 µg/mL, Immunohistochemistry: 5-20 µg/mL, Immunocytochemistry: 5-20 µg/mL, Optimal working dilutions must be determined by end user.

Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 500 µg/mL

Buffer: 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.