

Datasheet for ABIN7634673

anti-Angiotensin II antibody



Overview

Quantity:	100 μL
Target:	Angiotensin II (Ang II)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Angiotensin II antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Monoclonal Antibody to Angiotensin II (AngII)
Immunogen:	CPA005Hu210VA Conjugated Angiotensin II (AngII)
Clone:	C7
Specificity:	The antibody is a mouse monoclonal antibody raised against Angll. It has been selected for its
	ability to recognize Angll in immunohistochemical staining and western blotting.
Purification:	Protein A + Protein G affinity chromatography

Target Details

Target:	Angiotensin II (Ang II)
Alternative Name:	Angli (Ang Il Products)

Target Details

Target Details		
Background:	Ang-II, Angiotensin-2	
Pathways:	JAK-STAT Signaling, ACE Inhibitor Pathway, EGFR Signaling Pathway, Peptide Hormone	
	Metabolism, Regulation of Systemic Arterial Blood Pressure by Hormones, Regulation of Lipid	
	Metabolism by PPARalpha, Protein targeting to Nucleus, Feeding Behaviour, Regulation of long-	
	term Neuronal Synaptic Plasticity	
Application Details		
Application Notes:	Western blotting: 0.5-2 μg/mL,Immunohistochemistry: 5-20 μg/mL,Immunocytochemistry: 5-	
	20 μg/mL,Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated	
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious	
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration	
	date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	
Preservative:	ProClin, Sodium azide	
Precaution of Use:	This product contains ProClin and Sodium azide: POISONOUS AND HAZARDOUS SUBSTANCES	
	which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without	
	detectable loss of activity. Avoid repeated freeze-thaw cycles.	