

Datasheet for ABIN7635545

anti-BCAT2 antibody



Overview

Quantity:	100 μL
Target:	BCAT2
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This BCAT2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Troduct Details	
Purpose:	Polyclonal Antibody to Branched Chain Aminotransferase 2, Mitochondrial (BCAT2)
Immunogen:	RPD859Hu01Recombinant Branched Chain Aminotransferase 2, Mitochondrial (BCAT2)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against BCAT2. It has been selected for its ability to recognize BCAT2 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Pig
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	
Target:	BCAT2

Target Details

Target Details	
Alternative Name:	BCAT2 (BCAT2 Products)
Background:	BCAM, BCT2, PP18, BCATM, ECA40, Placental protein 18, Branched-chain-amino-acid aminotransferase, mitochondrial
UniProt:	015382
Application Details	
Application Notes:	Western blotting: 0.2-2 μ g/mL,1:250-2500 Immunohistochemistry: 5-20 μ g/mL,1:25-100 Immunocytochemistry: 5-20 μ g/mL,1:25-100 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.