

Datasheet for ABIN7635918 **anti-CEACAM6 antibody**



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	CEACAM6
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CEACAM6 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Carcinoembryonic Antigen Related Cell Adhesion Molecule 6 (CEACAM6)
Immunogen:	RPC980Hu02Recombinant Carcinoembryonic Antigen Related Cell Adhesion Molecule 6 (CEACAM6)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against CEACAM6. It has been selected for its ability to recognize CEACAM6 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	CEACAM6
---------	---------

Target Details

Alternative Name:	CEACAM6 (CEACAM6 Products)
Background:	CD66c, NCA, CD66-C, CEAL, Non-Specific Cross Reacting Antigen, Normal cross-reacting antigen
UniProt:	P40199

Application Details

Application Notes:	Western blotting: 0.01-2 µg/mL, Immunohistochemistry: 5-20 µg/mL, Immunocytochemistry: 5-20 µg/mL, Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.