

## Datasheet for ABIN7636195

## anti-CEBPA antibody



|  |  | oo to i roddot pago |
|--|--|---------------------|
|  |  |                     |
|  |  |                     |
|  |  |                     |

| ( | )             | V  |          | rV | ĺ | 9             | V | V |
|---|---------------|----|----------|----|---|---------------|---|---|
| ' | $\mathcal{I}$ | ٧V | <u> </u> | v  | 1 | $\overline{}$ | ٧ | ٧ |

| Quantity:    | 100 μL   |
|--------------|--|
| Target:      | CEBPA  |
| Reactivity:  | Rat  |
| Host:        | Mouse  |
| Clonality:   | Monoclonal   |
| Conjugate:   | This CEBPA antibody is un-conjugated   |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunoprecipitation (IP) |

## **Product Details**

| Purpose:       | Monoclonal Antibody to CCAAT/Enhancer Binding Protein Alpha (CEBPa)   |
|----------------|---|
| Immunogen:     | RPA644Ra01Recombinant CCAAT/Enhancer Binding Protein Alpha (CEBPa)  |
| Clone:         | C9  |
| Specificity:   | The antibody is a mouse monoclonal antibody raised against CEBPa. It has been selected for its ability to recognize CEBPa in immunohistochemical staining and western blotting. |
| Purification:  | Protein A + Protein G affinity chromatography   |
| Target Details |   |

| Target:           | CEBPA                  |
|-------------------|------------------------|
| Alternative Name: | CEBPa (CEBPA Products) |

## **Target Details**

| raiget Details      |   |
|---------------------|---|
| Background:         | CEBP-A  |
| UniProt:            | P05554  |
| Pathways:           | Brown Fat Cell Differentiation, Positive Regulation of fat Cell Differentiation   |
| Application Details |   |
| Application Notes:  | Western blotting: 0.5-2 μg/mL,Immunohistochemistry: 5-20 μg/mL,Immunocytochemistry: 5-20 μg/mL,Optimal working dilutions must be determined by end user.  |
| Comment:            | The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition. |
| Restrictions:       | For Research Use only   |
| Handling            |   |
| Format:             | Liquid  |
| Concentration:      | 1 mg/mL   |
| Buffer:             | 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.  |
| Preservative:       | ProClin   |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.   |
| Storage:            | 4 °C,-20 °C   |
| Storage Comment:    | Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without   |

detectable loss of activity. Avoid repeated freeze-thaw cycles.