

Datasheet for ABIN7636750

anti-CXCL5 antibody



Overview

Quantity:	100 μL
Target:	CXCL5
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CXCL5 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Monoclonal Antibody to Epithelial Neutrophil Activating Peptide 78 (ENA78)
Immunogen:	RPA860Hu02Recombinant Epithelial Neutrophil Activating Peptide 78 (ENA78)
Clone:	1#
Specificity:	The antibody is a mouse monoclonal antibody raised against ENA78. It has been selected for its ability to recognize ENA78 in immunohistochemical staining and western blotting.
Purification:	Protein A + Protein G affinity chromatography
Target Details	

Target Details

Target:	CXCL5
Alternative Name:	ENA78 (CXCL5 Products)

Target Details	
Background:	CXCL5, SCYB5, Chemokine C-X-C-Motif Ligand 5, Small Inducible Cytokine Subfamily B(Cys-X-Cys),Member 5, Neutrophil-activating peptide ENA-78
UniProt:	P42830
Pathways:	Cellular Response to Molecule of Bacterial Origin, Regulation of Leukocyte Mediated Immunity
Application Details	
Application Notes:	Western blotting: 0.01-2 μ g/mL,lmmunohistochemistry: 5-20 μ g/mL,lmmunocytochemistry: 5-20 μ g/mL,Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.